

**VARIETIES OF GROUPOIDS WITH AXIOMS OF THE  
FORM  $x^{m+1}y = xy$  AND/OR  $xy^{n+1} = xy$**

ĀORĀI ČUPONA, NAUM CELAKOSKI AND BILJANA JANEVA  
Macedonian Academy of Sciences and Arts and Ss. Cyril and Methodius  
University, Skopje, R. Macedonia

ABSTRACT. The subject of this paper are varieties  $\mathcal{U}(M; N)$  of groupoids defined by the following system of identities

$$\{x^{m+1} \cdot y = xy : m \in M\} \cup \{x \cdot y^{n+1} = xy : n \in N\},$$

where  $M, N$  are sets of positive integers. The equation  $\mathcal{U}(M; N) = \mathcal{U}(M'; N')$  for any given pair  $(M, N)$  is solved, and, among all solutions, one called canonical, is singled out. Applying a result of Evans ([6]) it is shown for finite  $M$  and  $N$  that: if  $M$  and  $N$  are nonempty and  $\gcd(M) = \gcd(M \cup N)$ , or only one of  $M$  and  $N$  is nonempty, then the word problem is solvable in  $\mathcal{U}(M; N)$ .

1. INTRODUCTION

A *groupoid* is an algebra  $\mathbf{G} = (G, \bullet)$  with one binary operation  $\bullet : (a, b) \mapsto ab$ . (We will often omit the operation sign.) Assuming the usual meanings of other algebraic notions, we do not define them explicitly.

By a result of P. Hall (see, for example, [3], III.2, Ex. 2, p. 125, or [10], p. 39-40), for any positive integer  $k$  there exist  $\frac{(2k-2)!}{k!(k-1)!}$  *k*-th *groupoid powers*  $x \mapsto x^k$ . In this paper, we assume the groupoid power  $x^k$  defined as follows:

$$x^1 = x, \quad x^{k+1} = x^k x.$$

So  $x^3 = x^2 x = (xx)x$ .

A formula  $x^{k+1}y = xy$  ( $xy^{k+1} = xy$ ), will be called a *left (right) equation*. (Here, and further on,  $m, n, k, p, i, j, s$  are assumed to be positive integers, and  $xy^{n+1}$  stands for  $x \cdot y^{n+1}$ , and  $x^{m+1}y$  for  $x^{m+1} \cdot y$ .) The varieties

---

2000 *Mathematics Subject Classification*. 03C05, 03D40, 08A50, 08A55, 08B20.

*Key words and phrases*. Groupoids, varieties of groupoids, partial groupoids, free groupoids, word problem.

$\mathcal{U}(M; \emptyset)$ ,  $\mathcal{U}(\emptyset; N)$ ,  $\mathcal{U}(M; N)$ , where  $M \neq \emptyset$  and  $N \neq \emptyset$ , are said to be *left*, *right*, *two-sided*, respectively. (Throughout the paper "variety" will mean "left, right, or two-sided variety".)

Below,  $\mathcal{U}(m_1, m_2, m_3, \dots; n_1, n_2, n_3, \dots)$  will be an abbreviation for  $\mathcal{U}(\{m_1, m_2, m_3, \dots\}; \{n_1, n_2, n_3, \dots\})$ .

The paper consists of three sections. In Section 2 we show that each variety  $\mathcal{U}(M; N)$  admits a canonical axiom system. In Section 3 we solve the equation  $\mathcal{U}(M; N) = \mathcal{U}(M'; N')$ . Finally, in Section 4, we consider "incomplete  $\mathcal{U}(M; N)$ -groupoids", and applying a result of Evans ([6]) we show that the word problem is solvable in  $\mathcal{U}(M; N)$  for finite  $M$  and  $N$  in each of the cases: (i)  $N = \emptyset$ , (ii)  $M = \emptyset$ , (iii)  $M \neq \emptyset$ ,  $N \neq \emptyset$ ,  $\gcd(M) = \gcd(M \cup N)$ .<sup>1</sup>

## 2. A CANONICAL AXIOM SYSTEM FOR $\mathcal{U}(M; N)$

The main result of this section is the following

THEOREM 2.1. *If  $M, N$  are nonempty sets of positive integers, then*

- (l)  $\mathcal{U}(M; \emptyset) = \mathcal{U}(\gcd(M); \emptyset)$ .
- (r)  $\mathcal{U}(\emptyset; N) = \mathcal{U}(\emptyset; \langle N \rangle)$ .<sup>2</sup>
- (t)  $\mathcal{U}(M; N) = \mathcal{U}(\gcd(M); \gcd(M \cup N))$ .

In order to prove this theorem we will show some lemmas, where  $m, n, k, p, i, j, s$  are assumed to be positive integers as above, and  $q$  a non-negative integer.

LEMMA 2.2. *If  $1 \leq k \leq m$ , then*

$$\mathcal{U}(m; \emptyset) \models x^{qm+k+1} = x^{k+1}. \quad 3$$

PROOF. Clearly,  $x^{m+2} = x^2, \dots, x^{2m+1} = x^{m+1}$  are true in  $\mathcal{U}(m; \emptyset)$ ; then the proof follows by induction on  $q$  and  $k$ .  $\square$

As a corollary, we obtain:

LEMMA 2.3. *If  $m|n$ , then  $\mathcal{U}(m; \emptyset) \subseteq \mathcal{U}(n; \emptyset)$ .*<sup>4</sup>

LEMMA 2.4. *If  $\gcd(M) = d \notin M$ , then there exists a nonempty set  $M_1$  of positive integers such that*

$$(2.1) \quad \mathcal{U}(M; \emptyset) = \mathcal{U}(M_1; \emptyset), \quad d = \gcd(M_1), \quad \min(M_1) < \min(M). \quad 5$$

PROOF. Let  $p = \min(M)$ . The assumption  $d \notin M$  implies that  $d < p$  and thus there exists an  $n \in M$  such that  $p$  is not a divisor of  $n$ . Then  $n = qp + k$ ,  $d|k$ ,  $k < p$  and, if  $M_1 = (M \setminus \{n\}) \cup \{k\}$ , the relations (2.1) hold.  $\square$

<sup>1</sup> $\gcd(M)$  is the greatest common divisor of  $M$

<sup>2</sup> $\langle N \rangle$  is the additive groupoid of integers generated by  $N$ .

<sup>3</sup> $\mathcal{V} \models \tau_1 = \tau_2$  means: the equation  $\tau_1 = \tau_2$  is true in the variety  $\mathcal{V}$ .

<sup>4</sup> $m|n$  denotes that  $m$  is a divisor of  $n$ .

<sup>5</sup> $\min(M)$  denotes the least element in  $M$ .

As a corollary of Lemma 2.3 and Lemma 2.4 we obtain the equality (l). The equality (r) is an obvious corollary of the following

LEMMA 2.5.  $\mathcal{U}(\emptyset; m, n) \subseteq \mathcal{U}(\emptyset; m + n)$ .

PROOF.  $\mathcal{U}(\emptyset; m) \models (x^{m+1})^i = x^{m+i}$ , and therefore  $\mathcal{U}(\emptyset; m, n) \models (x^{m+1})^{n+1} = x^{m+n+1}$ . Thus, if  $\mathbf{G} \in \mathcal{U}(\emptyset; m, n)$ , then:

$$x^{m+n+1}y = (x^{m+1})^{n+1} = x^{m+1}y = xy, \text{ i.e. } \mathbf{G} \in \mathcal{U}(\emptyset; m + n).$$

□

It remains to prove (t).

LEMMA 2.6. *If  $L = \{\gcd(m, n) : n \in N\}$ , then  $\mathcal{U}(m; N) = \mathcal{U}(m; L)$ .*

PROOF. By a similar argument as in Lemma 2.3,  $\mathcal{U}(m; L) \subseteq \mathcal{U}(m; N)$ . If  $n \in N$  and  $d = \gcd(m, n)$ , then there exist  $i, j$  such that  $im + d = jn$ . By Lemma 2.2,  $\mathcal{U}(m; n) \models x^{d+1} = x^{im+d+1}$ , and therefore  $\mathcal{U}(m; n) \models xy^{d+1} = xy$ . □

In completing the proof of (t) we will use the following result (for example [5] or [9]).

LEMMA 2.7. *If  $S$  is an additive groupoid of positive integers and  $d = \gcd(S)$ , then:*

- (i)  $\gcd(N) = d$  for any generating subset  $N$  of  $S$ .
- (ii) *There exists the least generating subset  $K = \{n_1, n_2, \dots, n_k\}$  of  $S$ , and  $K$  is finite.*
- (iii) *There exists  $s \in S$  such that for each positive integer  $j$ ,  $s + jd \in S$ .*

LEMMA 2.8. *If  $d_1, d_2, \dots, d_k$  are divisors of  $m$  and  $d = \gcd(d_1, d_2, \dots, d_k)$ , then*

$$\mathcal{U}(m; d_1, d_2, \dots, d_k) = \mathcal{U}(m; d).$$

PROOF. The inclusion  $\mathcal{U}(m; d) \subseteq \mathcal{U}(m; d_1, d_2, \dots, d_k)$  follows as in Lemma 2.6. For the converse inclusion, denote by  $S$  the additive groupoid of positive integers generated by  $\{d_1, d_2, \dots, d_k\}$ . By Lemma 2.7 (i) and (r) we have  $\gcd(S) = d$ , and  $\mathcal{U}(m; d_1, d_2, \dots, d_k) = \mathcal{U}(m; S)$ . By Lemma 2.7 (iii) there exists  $s \in S$  such that  $ms + d \in S$  and thus, by Lemma 2.2,  $\mathcal{U}(m; S) \models y^{ms+d+1} = y^{d+1}$ . □

Finally, by (l), (r), Lemma 2.6 and Lemma 2.8, it follows that

$$\mathcal{U}(M; N) = \mathcal{U}(m; n),$$

where  $m = \gcd(M)$  and  $n = \gcd(M \cup N)$ . This completes the proof of (t).

We note that the following equality holds in  $\mathcal{U}(m; m)$

$$(2.2) \quad (x^{m+1})^{m+1} = x^{m+1},$$

(or more generally, in  $\mathcal{U}(m; n)$ , where  $n|m$ , the equality  $(x^{in+1})^{m+1} = x^{in+1}$  holds.)

The results obtained in Theorem 2.1 suggest saying that

$$x^{m+1}y = xy, \{xy^{n+1} = xy : n \in K\}, \{x^{m+1}y = xy, xy^{n+1} = xy\}$$

is the *canonical axiom system* of  $\mathcal{U}(M; \emptyset)$ ,  $\mathcal{U}(\emptyset; N)$ ,  $\mathcal{U}(M; N)$ , respectively, where  $M, N$  are nonempty sets of positive integers,  $m = \gcd(M)$ ,  $K$  is the least generating subset of  $\langle N \rangle$ , and  $n = \gcd(M \cup N)$ .

As a corollary of Theorem 2.1 (for example [2]) we obtain

**COROLLARY 2.9.** *For any pair  $(M, N)$  the variety  $\mathcal{U}(M; N)$  is finitely based.*

### 3. CLOSED SETS OF EQUATIONS IN $\mathcal{U}(M; N)$

The main result of this section is the following

**THEOREM 3.1.** *If  $M, N, M', N'$  are nonempty sets of positive integers, then:*

- (i)  $\mathcal{U}(M; \emptyset) = \mathcal{U}(M'; \emptyset) \iff \gcd(M) = \gcd(M')$ .
- (ii)  $\mathcal{U}(\emptyset; N) = \mathcal{U}(\emptyset; N') \iff \langle N \rangle = \langle N' \rangle$ .
- (iii)  $\mathcal{U}(M; N) = \mathcal{U}(M'; N') \iff$   
 $\gcd(M) = \gcd(M') \ \& \ \gcd(M \cup N) = \gcd(M' \cup N')$ .
- (iv)  $\mathcal{U}(M; \emptyset) \neq \mathcal{U}(\emptyset; N)$ ;  $\mathcal{U}(M; \emptyset) \neq \mathcal{U}(M'; N')$ ;  $\mathcal{U}(\emptyset; N) \neq \mathcal{U}(M'; N')$ .

The  $\Leftarrow$ -parts of (i), (ii), (iii) hold by Theorem 2.1. The corresponding  $\Rightarrow$ -parts and (iv) are corollaries of the following statement, shown in [4] (Proposition 3.5).

**PROPOSITION 3.2.** *Let  $\mathbf{H}$  be a free groupoid in the variety  $\mathcal{U}(M; N)$ . Then the following statements hold:*

- (i) *If  $M \neq \emptyset$ ,  $N = \emptyset$ ,  $\gcd(M) = m$ , then a left equation  $x^{n+1}y = xy$  holds in  $\mathbf{H}$  iff  $m|n$ ; no right equation holds in  $\mathbf{H}$ .*
- (ii) *If  $M = \emptyset$ ,  $N \neq \emptyset$ , then a right equation  $xy^{n+1} = xy$  holds in  $\mathbf{H}$  iff  $n \in \langle N \rangle$ ; no left equation holds in  $\mathbf{H}$ .*
- (iii) *If  $M \neq \emptyset$ ,  $N \neq \emptyset$  and  $m = \gcd(M)$ ,  $n = \gcd(M \cup N)$ , then  $x^{i+1}y = xy$  iff  $m|i$ , and  $xy^{j+1} = xy$  iff  $n|j$ , hold in  $\mathbf{H}$ .*

(We note that only-if parts of (i) and (iii) in Proposition 3.2 follow from the fact that  $C_n \in \mathcal{U}(n; \emptyset) \cap \mathcal{U}(kn; n)$ , where  $C_n$  is the groupoid that is the reduction of the cyclic group of order  $n$  to its binary operation.)

A set  $\Sigma$  of equations is said to be *closed* if, for every equation  $\varepsilon$ , the following implication holds:

$$(\Sigma \models \varepsilon) \Rightarrow (\varepsilon \in \Sigma).$$

PROPOSITION 3.3. (i) Assume that  $\Sigma$  is a set of equations containing at least one left equation and at least one right equation. Then  $\Sigma$  is a closed set iff there exist two positive integers  $m$  and  $n$  such that  $n$  is a divisor of  $m$  and

$$\Sigma = \{x^{im+1}y = xy : i \geq 1\} \cup \{xy^{jn+1} = xy : j \geq 1\}.$$

(ii) A set  $\Sigma$  of left equations is closed iff there is a positive integer  $m$  such that

$$\Sigma = \{x^{im+1}y = xy : i \geq 1\}.$$

(iii) A set  $\Sigma$  of right equations is closed iff there is an additive groupoid  $S$  of positive integers such that

$$\Sigma = \{xy^{n+1} = xy : n \in S\}.$$

The lattices  $\mathcal{U}_l, \mathcal{U}_r, \mathcal{U}$  (of all left, right, two-sided varieties, respectively) can be characterized as follows:

PROPOSITION 3.4. (l)  $\mathcal{U}_l$  is isomorphic to the lattice of positive integers, where  $m \leq n$  iff  $m|n$ .

(r)  $\mathcal{U}_r$  is antiisomorphic to the lattice of additive groupoids of positive integers.

(t)  $\mathcal{U}$  is isomorphic to the lattice of pairs  $(m, n)$  of positive integers such that  $n$  is divisor of  $m$ , and:

$$(m, n) \leq (m', n') \iff m|m' \ \& \ n|n'.$$

#### 4. INCOMPLETE $\mathcal{U}(M; N)$ - GROUPOIDS AND VARIETIES $\mathcal{U}(M; N)$ WITH SOLVABLE WORD PROBLEM

We investigate here the class of incomplete  $\mathcal{U}(M; N)$ -groupoids and by applying the main result of Evans's paper [6], we solve the word problem for some varieties  $\mathcal{U}(M; N)$ .

The term "incomplete groupoid" ([6]) has the same meaning as "half-groupoid" ([1]) or "partial groupoid" ([8]). Namely, if  $G$  is a nonempty set,  $D$  a subset of  $G \times G$ , and  $\cdot : (x, y) \mapsto xy$  a map from  $D$  into  $G$ , then the pair  $\mathbf{G} = (G, \cdot)$  is called an *incomplete groupoid* with the *domain*  $D$ .

A groupoid  $\mathbf{H} = (H, \bullet)$  is called an *extension* of the incomplete groupoid  $\mathbf{G}$  iff  $G \subseteq H$  and  $a \bullet b = ab$ , for every  $(a, b) \in D$ . If  $G^o = G \cup \{0\}$ , where  $0 \notin G$ , then the groupoid  $\mathbf{G}^o = (G^o, \bullet)$  defined by

$$(4.1) \quad x \bullet y = \begin{cases} xy, & \text{if } (x, y) \in D \\ 0, & \text{otherwise} \end{cases}$$

is an extension of  $\mathbf{G}$ . We call  $\mathbf{G}^o$  the *trivial extension* of  $\mathbf{G}$ .

If  $M, N$  are sets of positive integers such that  $M \cup N \neq \emptyset$ , then we denote by  $\mathcal{IU}(M; N)$  the *class of incomplete groupoids*  $\mathbf{G}$ , such that the corresponding

trivial closure  $\mathbf{G}^o$  satisfies the following implications:

$$(4.2) \quad \begin{aligned} x^{m+1} \in G &\Rightarrow x^{m+1} \bullet y = x \bullet y, \\ y^{n+1} \in G &\Rightarrow x \bullet y^{n+1} = x \bullet y, \end{aligned}$$

for any  $m \in M, n \in N, x, y \in G$ .

Let  $\mathbf{G}$  be an incomplete groupoid and  $K$  a set of positive integers. We define an *equivalence*  $\sim_K$  on  $G$  as follows. If  $K = \emptyset$ , then  $\sim_K$  is the equality on  $G$ . If  $K \neq \emptyset$ , we define a relation  $\rightarrow_K$  on  $G$  by:

$$(4.3) \quad c \rightarrow_K d \iff d = c^{k+1},$$

for  $c, d \in G$  and some  $k \in K$ , and we put:  $c \leftrightarrow_K d \iff (c \rightarrow_K d \text{ or } c \leftarrow_K d)$ . We denote by  $\sim_K$  the reflexive, symmetric and transitive closure of  $\rightarrow_K$  on  $G$ , i.e., the equivalence on  $G$  generated by  $\rightarrow_K$ .

By (4.1), (4.2), and (4.3), we obtain the following characterization of the class  $\mathcal{IU}(M; N)$ :

$$(4.4) \quad \mathbf{G} \in \mathcal{IU}(M; N) \iff (\forall x, x', y, y' \in G)(x \sim_M x' \ \& \ y \sim_N y' \Rightarrow xy = x'y')$$

Let  $\mathbf{G} \in \mathcal{IU}(M; N)$  and define

$$(4.5) \quad A = \{a \in G \mid a^{k+1} \in G, \text{ for every } k \in M \cup N\}, \quad B = G \setminus A;$$

clearly,  $B = \{b \in G \mid b^{k+1} \notin G, \text{ for some } k \in M \cup N\}$ .

By (4.1), (4.2) and (4.5) it follows that

$$(4.6) \quad \mathbf{G} \in \mathcal{IU}(M; N) \ \& \ A = G \Rightarrow \mathbf{G}^o \in \mathcal{U}(M; N).$$

Note that, in the special case when  $M = \{m\}, N = \{n\}$ , and  $n|m$ , we have  $A = \{a \in G \mid a^{m+1} \in G\}$  and  $B = \{b \in G \mid b^{m+1} \notin G\}$ .

The following proposition is true.

- PROPOSITION 4.1. (i) If  $\mathbf{G} \in \mathcal{IU}(m; \emptyset)$ , then for each  $a \in A, q \geq 0$ , and  $1 \leq k \leq m$ , the equality  $a^{qm+k+1} = a^{k+1}$  holds.  
(ii) If  $\mathbf{G} \in \mathcal{IU}(m; n), n|m$ , and  $a \in A$ , then  $(a^{in+1})^{m+1} = a^{in+1}$ .  
(iii)  $\mathcal{IU}(\emptyset; r, i) = \mathcal{IU}(\emptyset; r, r+i)$ .

Using (4.3) and Proposition 4.1 we obtain the following

LEMMA 4.2. Let  $\mathbf{G} \in \mathcal{IU}(m; n)$  and  $n|m$ . Then

- (i)  $x \sim_m y \Rightarrow x^{m+1} = y^{m+1}$ ;  
(ii)  $x \sim_m y \Rightarrow x, y \in A \vee x = y \in B$ ,

where  $\sim_m$  stands for  $\sim_{\{m\}}$ .

PROOF. Let  $x \sim_m y$ . If  $x = y$ , then  $x^{m+1} = y^{m+1}$ . If  $x \neq y$ , then  $x \sim_m y \iff (\exists t_0, t_1, \dots, t_s \in G)x = t_0 \leftrightarrow t_1 \leftrightarrow \dots \leftrightarrow t_s = y$ , where  $\leftrightarrow$  stands for  $\leftrightarrow_{\{m\}}$ . The proof is given by induction on  $s$ . If  $s = 1$ , then  $x^{m+1} = y^{m+1}$ , and  $x, y \in A$ . If  $s = 2$ , we have the following four cases:

- 1)  $x \rightarrow t \rightarrow y$ ; then  $t = x^{m+1}$ ,  $y = t^{m+1}$ ,  $y = (x^{m+1})^{m+1} = x^{m+1}$   
(by Proposition 4.1), and thus  $y^{m+1} = (x^{m+1})^{m+1} = x^{m+1}$ ;
- 2)  $x \rightarrow t \leftarrow y$ ; then  $x^{m+1} = t = y^{m+1}$ ;
- 3)  $x \leftarrow t \leftarrow y$ ; then  $x^{m+1} = y^{m+1}$  follows by symmetry of 1);
- 4)  $x \leftarrow t \rightarrow y$ ; then  $x = t^{m+1} = y$ ;

and in each case  $x, y \in A$ .

If  $s > 2$ , then applying 1)–4), the sequence  $t_0, t_1, \dots, t_s$  can be reduced to a sequence with less than  $s + 1$  elements.  $\square$

As a corollary of Lemma 4.2 we obtain the following

PROPOSITION 4.3.

$$(4.7) \quad \mathbf{G} \in \mathcal{IU}(m; n) \ \& \ n|m \Rightarrow (\forall b, b' \in B)(b \sim_m b' \Rightarrow b = b').$$

If  $b \in B$ , then we denote by  $p(b)$  the positive integer  $p$ , such that

$$(4.8) \quad b^p \neq 0, \ b^{p+1} = 0.$$

Now we are ready to prove the main result.

THEOREM 4.4. *If the pair  $(M, N)$  satisfies one of the following conditions*

- (i)  $M = \emptyset$ ,  $N \neq \emptyset$ ; (ii)  $M = \{m\}$ ,  $N = \emptyset$ ; (iii)  $M = \{m\} = N$ ,

*then for each (finite)  $\mathbf{G} \in \mathcal{IU}(M; N)$  there exists a (finite)  $\mathbf{H} \in \mathcal{U}(M; N)$  that is an extension of  $\mathbf{G}$ .*

PROOF. If  $B = \emptyset$ , then  $\mathbf{G}^o$  is an extension of  $\mathbf{G}$ , finite if  $G$  is finite, such that, by (4.6),  $\mathbf{G}^o \in \mathcal{U}(M; N)$ . Thus, it remains to build an extension  $\mathbf{H} = (H, \bullet) \in \mathcal{U}(M; N)$ , assuming that  $B \neq \emptyset$ .

Consider first the case (i):  $M = \emptyset$ ,  $N \neq \emptyset$ .

Let  $L$  be a set such that  $L \cap G^o = \emptyset$ , and let  $b \mapsto \underline{b}$  be a surjection from  $B$  onto  $L$  with the following property:

$$(4.9) \quad (\forall b, c \in B)(\underline{b} = \underline{c} \iff b \sim c \ \& \ b^p = c^q),$$

where  $\sim$  is an abbreviation for  $\sim_N$ ,  $p = p(b)$ ,  $q = p(c)$ . Define an operation  $\bullet$  on  $H = G^o \cup L$  as follows:

- 1) If  $x, y \in G$ ,  $b \in B$ , then:
  - 1.1)  $x \bullet y = xy$ , for  $xy \in G$ ;
  - 1.2)  $x \bullet y = \underline{b}$ , for  $x = b^p, y \sim b$ .
- 2) If  $x \in G, b \in B$ , then:
  - 2.1)  $\underline{b} \bullet x = \underline{b}$ , for  $x \sim b$ ;
  - 2.2)  $x \bullet \underline{b} = x \bullet b$ , if  $x \bullet b$  is defined by 1.1) or 1.2).
- 3) If  $b, c \in B$ , and  $b \sim c$ , then  $\underline{b} \bullet \underline{c} = \underline{b}$ .
- 4)  $x \bullet y = 0$ , in any other case.

Using (4.9) and (4.4) one can directly show that  $\bullet$  is a well-defined operation on  $H$ .

It follows by 1.1) that  $\mathbf{H}$  is an extension of  $\mathbf{G}$ , and so it remains to show that  $\mathbf{H} \in \mathcal{U}(\emptyset; N)$ .

First, by (4.9) and the definition of  $\bullet$  we obtain the following properties:

- 5) If  $a \in A$ ,  $b \in B$ ,  $z \in L \cup \{0\}$ ,  $n \in N$ ,  $p = p(b)$ , then:
- 5.1)  $a_{\bullet}^{n+1} = a^{n+1}$ ;
  - 5.2)  $b_{\bullet}^{n+1} = b^{n+1}$ , for  $n + 1 \leq p$ ;
  - 5.3)  $b_{\bullet}^{n+1} = \underline{b}$ , for  $n + 1 > p$ ;
  - 5.4)  $z_{\bullet}^k = z$ , for each  $k \in Z^+$ .

(Here,  $y_{\bullet}^k$  is the  $k$ -th power of  $y$  in  $\mathbf{H}$ , i.e.  $y_{\bullet}^1 = y$ ,  $y_{\bullet}^{k+1} = y_{\bullet}^k \bullet y$ .)

Now, by using properties 5) and the definition of  $\bullet$ , we can show that:

- 6)  $x \bullet (y_{\bullet}^{n+1}) = x \bullet y$ , for each  $x, y \in H$ ,  $n \in N$ , i.e.  $\mathbf{H} \in \mathcal{U}(\emptyset; N)$ .

Thus we have proved Theorem 4.4 in the case (i).

Now, consider the cases (ii)  $M = \{m\}$ ,  $N = \emptyset$  and (iii)  $M = N = \{m\}$ . The construction of a groupoid  $\mathbf{H} \in \mathcal{U}(M; N)$  that is an extension of  $\mathbf{G} \in \mathcal{IU}(M; N)$  is formally the same in case (ii) as in case (iii). In both cases we will denote the equivalence  $\sim_M$  in  $G$  by  $\sim$ ; and  $\approx$  is the equality in  $G$  in case (ii), and  $\approx$  is the same as  $\sim$  in case (iii).

Let

$$L = \{(b, i) : b \in B, p(b) < i \leq m\}$$

and  $H = G^o \cup L$ . (The union defining  $H$  is assumed to be disjoint.)

Define an operation  $\bullet$  in  $H$  as follows.

- 1') If  $x, y \in G$ , then:
- 1.1')  $x \bullet y = xy$ , if  $xy \in G$ ;
  - 1.2')  $x \bullet y = b$ , if  $b \in B$ ,  $x \sim b^m$ ,  $p(b) = m$ ,  $y \approx b$ ;
  - 1.3')  $x \bullet y = (b, p(b) + 1)$ , if  $x \sim b^{p(b)}$ ,  $p(b) < m$ ,  $y \approx b$ .
- 2') If  $b \in B$ ,  $y \in G$ ,  $y \approx b$ , then:
- 2.1')  $(b, m) \bullet y = b$ ;
  - 2.2')  $(b, i) \bullet y = (b, i + 1)$ , if  $p(b) < i < m$ .
- 3') If  $x \in L$ , then  $x \bullet x = x$ .
- 4')  $x \bullet y = 0$ , in any other case.

Thus we obtain an extension  $\mathbf{H} = (H, \bullet)$  of  $\mathbf{G}$ . (The product  $x \bullet y$  for (ii) in the cases 1.2') and 1.3') is well-defined by (4.7).)

It remains to show that  $\mathbf{H} \in \mathcal{U}(M; N)$ .

For that purpose, note first that the following statements hold.

- 5') If  $a \in A$ ,  $x \in B \cup L \cup \{0\}$ , then
- 5.1')  $a_{\bullet}^{m+1} = a^{m+1} \in G$ ;
  - 5.2')  $x_{\bullet}^{m+1} = x$ .

(Here, as in 5),  $y_{\bullet}^k$  is the  $k$ -th power of  $y$  in  $\mathbf{H}$ .)

We will now show that:



6')  $x \bullet^{m+1} \bullet y = x \bullet y$ , for any  $x, y \in H$ .

Namely, if  $x \in B \cup L \cup \{0\}$  or  $y \in L \cup \{0\}$ , then the equality 6') follows from 3'), 4') and 5.2'). There remains the case  $x \in A$ ,  $y \in G$ . Here, by 5.1') and the definition 1.1'), 1.2'), 1.3') and 4'), we obtain the desired equality 6').

Hence (in the case  $M = \{m\}$ ,  $N = \emptyset$ ),  $\mathbf{H} \in \mathcal{U}(m; \emptyset)$ .

It remains to show that, for  $M = N = \{m\}$ , the following identity holds in  $\mathbf{H}$ :

7')  $x \bullet (y \bullet^{m+1}) = x \bullet y$ .

By the same reasoning as for 6'), the equality 7') is true whenever  $y \in B \cup L \cup \{0\}$  or  $x = 0$ . For  $x \in G \cup L$  and  $y \in A$ , one can show that 7') is also true, in the same way as for 6').

Hence (in the case  $M = N = \{m\}$ ),  $\mathbf{H} \in \mathcal{U}(m; m)$ , and this completes the proof of Theorem 4.4.  $\square$

The following statement is a special case of the main result of the paper [6]:

PROPOSITION 4.5. *If the pair  $(M, N)$  is such that for every  $\mathbf{G} \in \mathcal{IU}(M; N)$  there exists an extension  $\mathbf{H} \in \mathcal{U}(M; N)$ , then the word problem is solvable in the variety  $\mathcal{U}(M; N)$ .*

As a corollary of Theorem 2.1, Proposition 4.5 and Theorem 4.4, we obtain the following

THEOREM 4.6. *If  $M \cup N$  is finite and one of the following conditions holds:*

(i)  $N = \emptyset$ ; (ii)  $M \neq \emptyset, N \neq \emptyset$ , and  $\gcd(M) = \gcd(M \cup N)$ ; (iii)  $M = \emptyset$ ,  
*then the word problem is solvable in the variety  $\mathcal{U}(M; N)$ .*

REMARK 4.7. Theorem 2.1 and Theorem 3.1 suggest the following two questions:

a) Is the implication

$$\mathcal{U}(M; N) = \mathcal{U}(M'; N') \Rightarrow \mathcal{IU}(M; N) = \mathcal{IU}(M'; N')$$

true?

b) Is it true that, for every pair  $(M, N)$ , every  $\mathbf{G} \in \mathcal{IU}(M; N)$  has an extension  $\mathbf{H} \in \mathcal{U}(M; N)$ ?

The answer to both questions, in general, is negative, as the following example shows.

Let  $M$  be a nonempty set of positive integers,  $\gcd(M) = m$  and  $G = \{1, 2, \dots, m+1, m+2\}$ . Let  $\mathbf{G} = (G, \bullet)$  be an incomplete groupoid such that the corresponding canonical extension  $\mathbf{G}^\circ$  is defined as follows:

- $a_1)$   $i \bullet 1 = i + 1$ , if  $i = 1, 2, \dots, m + 1$ ;
- $a_2)$   $1 \bullet (m + 2) = 1$ ;

$$a_3) (m+1) \bullet (m+2) = m+1;$$

$$a_4) x \bullet y = 0, \text{ otherwise.}$$

If  $m \notin M$  and  $p = \min(M) > m+1$ , then  $x^{m+1} = 0$  for every  $x \in G$ ,  $n \in M$ , and thus, by (4.3),  $\mathbf{G} \in \mathcal{IU}(M; \emptyset)$ . On the other hand, we have  $1_{\bullet}^{m+1} \bullet 1 = (m+1) \bullet 1 = m+2 \neq 2 = 1 \bullet 1$ , which implies that  $\mathbf{G} \notin \mathcal{IU}(m; \emptyset)$ . Hence,  $\mathcal{IU}(m; \emptyset) \not\subseteq \mathcal{IU}(M; \emptyset)$ , i.e. the answer to the question *a*) is negative.

Also,  $\mathbf{G} \in \mathcal{IU}(M; \emptyset)$  cannot be embedded in an  $\mathbf{H} \in \mathcal{U}(M; \emptyset) (= \mathcal{U}(m; \emptyset))$ , because  $(1_{\bullet}^{m+1}) \bullet 1 = m+2 \neq 2 = 1 \bullet 1$ .

REMARK 4.8. Theorem 4.4 and the main result of [7] imply that, for each of the cases: i)  $M \neq \emptyset$ ,  $N = \emptyset$ ; ii)  $M \neq \emptyset \neq N$ ,  $\gcd(M) = \gcd(M \cup N)$ ; iii)  $M = \emptyset$ ,  $N \neq \emptyset$ , the embeddability problem: "For a finite  $\mathbf{G} \in \mathcal{IU}(M; N)$ , is there an extension  $\mathbf{H} \in \mathcal{U}(M; N)$ ?" is solvable.

REMARK 4.9. In connection with Theorem 4.6, the authors conjecture that, applying the main result of [7], one can obtain the following variant of Theorem 4.6: "If  $M \cup N$  is finite, then the word problem is solvable in  $\mathcal{U}(M; N)$ ."

#### REFERENCES

- [1] R.H. Bruck: A Survey of Binary Systems, Berlin–Göttingen–Heidelberg, 1956.
- [2] S. Burris and H.P. Sankappanavar: A Course in Universal Algebra, Springer–Verlag, Grad. Texts in Math., New York–Berlin, 1981.
- [3] P.M. Cohn: Universal Algebra, Harper's Series in Modern Math., 1965.
- [4] Ā. Āupona, N. Celakoski and B. Janeva: *Free Groupoids with Axioms of the Form  $x^{m+1}y = xy$  and/or  $xy^{n+1} = xy$* , Novi Sad J. Math. **29** No. 2, (1999), 131-147, Proc. VIII Int. Conf. "Algebra & Logic" (Novi Sad, 1998).
- [5] D. Dimovski: *Semigroups of Integers with Addition* (in Macedonian), Maced. Acad. Sci. and Arts, Contributions, IX 2–Nat. Sci and Math. (1977), 21–26.
- [6] T. Evans: *The Word Problem for Abstract Algebras*, J. London Math. Soc. **26** (1951), 64–71.
- [7] T. Evans: *Embeddability and the Word Problem*, J. London Math. Soc. **28** (1952), 76–80.
- [8] G. Grätzer: Universal Algebra, D. Van Nostrand Co., 1968.
- [9] A.I. Mal'cev: *Algoritmi i Rekursivnie funkcii* (in Russian), Moskva 1965.
- [10] S. Markovski: *Finite Mathematics* (in Macedonian), Skopje 1993.

Macedonian Academy of Sciences and Arts  
1000 Skopje, R. Macedonia

Faculty of Mechanical Engineering  
1000 Skopje, R. Macedonia

Faculty of Natural Sciences and Mathematics  
Institute of informatics  
PB 162, 1000 Skopje, R. Macedonia  
E-mail address: biljana@pmf.ukim.edu.mk

Received: 26.07.2000.

Revised: 21.11.2001.