

Distribution of demersal crustaceans in the southern Adriatic Sea

Nicola UNGARO¹, Chiara Alessandra MARANO², Luca CERIOLA¹
and Michele MARTINO²

¹*Laboratorio Provinciale di Biologia Marina, Molo Pizzoli (Porto), 70123 Bari, Italy*

²*Dipartimento di Produzione Animale, Facoltà di Agraria, Università degli Studi Via Amendola 126/a, 70126 Bari, Italy*

Occurrence, distribution, and assemblage of crustaceans in the southern Adriatic Sea were determined from data collected during five seasonal (summer) trawl surveys carried out during 1996-2000. Trawls were conducted at a bathymetric range of 10-800 m, by the same vessel and same sampling gear in the framework of the E.U. Project, MEDITS. Fifty-two benthic-epibenthic, and nektonic species (three stomatopods, 49 decapods) were caught. Species density data (individuals per km²) were processed according to multivariate techniques to describe the composition and distribution of main species assemblages. The assemblages were consistent throughout the surveyed area with some differences between the western and eastern zones, mostly those found on shelf bottoms. The influence of oceanographic features, other than depth, is discussed.

Key words: decapod crustaceans, stomatopods, demersal assemblages, multivariate analysis, Adriatic Sea

INTRODUCTION

Species distribution and assemblages have been studied in several areas of the Mediterranean Sea to explain the space-time distribution patterns of demersal species versus environmental characteristics (depth, substratum, etc.) and anthropic impacts (fisheries, etc.). Until 1996, information on assemblages of demersal species was reported only for the western side of the Adriatic (UNGARO *et al.*, 1995, 1998). Since 1996, an annual experimental trawl survey (part of the Mediterranean International Trawl Survey,

MEDITS) was conducted in the late spring-early summer in the central and western Mediterranean Basin. The same sampling scheme, vessel, and gear were used during the survey in the southern Adriatic between the Italian (south-western Adriatic) and Albanian (south-eastern Adriatic) sides (BERTRAND *et al.*, 1997). Therefore, information on demersal species in the entire southern Adriatic is now available (UNGARO *et al.*, 1999; UNGARO & MARANO, 2001).

Crustaceans are an important fraction of the total catch in the southern Adriatic (PETRUZZI *et al.* 1988; PASTORELLI *et al.*, 1996; MARSAN *et*

al., 2000) and Mediterranean (RELINI-ORSI & RELINI, 1972; ABELLÓ *et al.*, 1988; RIGHINI & AUTERI, 1989; CARTES & SARDÀ, 1992; MURA & CAU, 1992; PIPITONE & TUMBIOLO, 1993; TURSI *et al.*, 1993; RINELLI *et al.*, 1998; SPANÒ, 1998). Some of the species are valuable for fishery exploitation (e.g., *Nephrops norvegicus*, *Parapenaeus longirostris*, *Aristeus antennatus*, *Aristaeomorpha foliacea*) and therefore require assessment and management (RELINI *et al.*, 1999). The links between the above-mentioned zoological group and the trophodynamics of the marine environment were postulated by several authors (ABELLÓ *et al.*, 1988; CARTES *et al.*, 1994; UNGARO *et al.*, 1999).

In the present work, data from the 1996-2000 MEDITS trawl surveys were processed to create a list of and quantify crustacean species on the trawlable bottoms of the southern Adriatic and to identify the composition and distribution of the main assemblages. The gathered information will be useful in defining crustacean biodiversity and local fish assemblages, important for the environmental management of the Mediterranean basin (CADDY, 1998).

MATERIAL AND METHODS

Crustacean samples came from trawl surveys conducted in an area of about 24000 km² in the southern Adriatic Sea (Fig. 1). A large deep trench in the middle of the surveyed area (1230 m deep - the deepest area in the Adriatic) roughly separated the trawlable bottoms of the western and eastern sides. The bottoms on both sides of the basin were trawled at 112 stations during the spring and summer of 1996-2000 on the shelf and the upper slope (16-588 m depth). One haul (60 square nautical miles) was sampled at each station during each survey (72 hauls in the western area, 40 hauls in the eastern area per year; 560 hauls during the 5-year survey) using an otter trawl net (length approx. 40 m, wing spread approx. 18 m) with 10 mm mesh at the cod end. During the first survey (1996), the sampling design was randomly stratified into five bathymetric strata: 10-50 m, 51-100 m, 101-200 m, 201-500 m, and 501-800 m (BERTRAND *et al.*, 1997, 2002). The same sampling stations were sampled in following years. The tow duration was one hour on slope bottoms and 0.5 h on shelf bottoms.




Fig. 1. The study area

All crustacean species collected by the net were identified according to ZARIQUIEY-ALVAREZ (1968) and FALCIAI & MINERVINI (1992) and a list per haul was recorded (true plankton species were excluded). Species abundance data were standardized to kg km⁻² and no. km⁻² by the swept-area method (SPARRE & VENEMA, 1992). Occurrence, biomass and density indices, and bathymetric ranges were calculated for each species. Species abundances per sampling station matrix were processed using multivariate techniques such as cluster analysis (hierarchical agglomerative clustering; group average; LEGENDRE & LEGENDRE, 1987) and multi-dimensional scaling (MDS; KRUSKAL, 1964). Before processing, the data were double-root transformed to avoid a strong influence by the most abundant species in the samples (CLARKE & GREEN, 1988). Non-metric MDS (ARDISSON *et al.*, 1990) and two-dimensional space ordination were performed. The similarity

among samples was evaluated with the BRAY-CURTIS coefficient, after excluding species with an abundance value below 1% from each sample to minimize 'noise' elaboration. Data analysis was carried out with PRIMER software (CARR, 1996). The same analysis was performed for the 'species mean abundance per depth stratum' matrices to summarize information according to sampling scheme (western and eastern areas; five depth strata) and determine differences between species assemblages and distributions per survey, depth, and area.

RESULTS

The five surveys produced 52 benthic or nektobenthic crustacean species (3 stomatopods and 49 decapods). Thirty zoological families were identified, with Pandalidae, Galatheididae, and Paguridae being the most represented (7, 4, and 4 species, respectively; Table 1).

Table 1. Crustacean species collected during MEDITS 1996-2000 surveys in the Southern Adriatic Sea: sampling area and geographic position of 112 hauls (72 and 40 samples on Western and Eastern trawable bottoms, respectively)

ORDER	FAMILY	SPECIES
	Squillidae	<i>Rissoides desmaresti</i> (Risso, 1816) <i>Rissoides pallidus</i> (Gesbrecht, 1910) <i>Squilla mantis</i> (Linnaeus, 1758)
ASTACIDEA	Nephropidae	<i>Homarus gammarus</i> (Linnaeus, 1758) <i>Nephrops norvegicus</i> (Linnaeus, 1758)
BRACHYRHYNCHA	Portunidae	<i>Bathynectes maravigna</i> (Prestandrea, 1839) <i>Liocarcinus depurator</i> (Linnaeus, 1758) <i>Macropipus tuberculatus</i> (Roux, 1830)
	Geryonidae	<i>Geryon longipes</i> A. Milne Edwards, 1881
	Goneplacidae	<i>Goneplax rhomboides</i> (Linnaeus, 1758)
	Xanthidae	<i>Monodaeus couchii</i> (Couch, 1851) <i>Philumnus hirtellus</i> , (Linnaeus, 1761) <i>Xantho pilipes</i> A. Milne Edwards, 1867
CARIDEA	Alpheidae	<i>Alpheus glaber</i> (Olivi, 1792)
	Pandalidae	<i>Chlorotocus crassicornis</i> (Costa, 1871) <i>Plesionika acanthonotus</i> (S.I. Smith, 1882) <i>Plesionika antigai</i> Zariquiey Alvarez, 1955 <i>Plesionika edwardsii</i> (Brandt, 1851)

Table 1. Cont'd

		<i>Plesionika gigliolii</i> (Senna, 1903)
		<i>Plesionika heterocarpus</i> (Costa, 1871)
		<i>Plesionika martia</i> (A. Milne Edw., 1883)
	Pasiphaeidae	<i>Pasiphaea multidentata</i> Esmark, 1866
		<i>Pasiphaea sivado</i> (Risso, 1816)
	Crangonidae	<i>Pontocaris lacazei</i> (Gourret, 1887)
		<i>Pontophilus spinosus</i> (Leach, 1815)
	Processidae	<i>Processa canaliculata</i> Leach, 1915
DROMIACEA	Homolidae	<i>Homola barbata</i> (Fabricius, 1793)
		<i>Paramola cuvieri</i> (Risso, 1816)
	Latreillidae	<i>Latreillia elegans</i> Roux, 1830
GALATHEIDEA	Galatheidae	<i>Munida intermedia</i> A.Mil.Edw.& Bouv., 1899
		<i>Munida iris</i> Zariquiey Alvarez, 1952
		<i>Munida perarmata</i> A.Mil.Edw.& Bouv., 1894
		<i>Munida rugosa</i> (Fabricius, 1775)
OXYRHYNCHA	Majidae	<i>Anamathia rissoana</i> (Roux, 1828)
		<i>Macropodia longipes</i> (A.Mil.Edw.& Bouv., 1899)
	Phartenopidae	<i>Parthenope macrocheles</i> (Herbst, 1790)
OXYSTOMATA	Calappidae	<i>Calappa granulata</i> (Linnaeus, 1767)
		<i>Calappa turkayana</i> n. sp.
	Dorippidae	<i>Medorippe lanata</i> (Linnaeus, 1767)
PAGURIDEA	Diogenidae	<i>Dardanus arrossor</i> (Herbst, 1796)
	Paguridae	<i>Pagurus alatus</i> Fabricius, 1775
		<i>Pagurus cuanensis</i> , Bell 1846
		<i>Pagurus excavatus</i> (Herbst, 1791)
		<i>Pagurus prideaux</i> Leach, 1815
PALINURIDEA	Palinuridae	<i>Palinurus elephas</i> (Fabricius, 1787)
	Polychelidae	<i>Polycheles typhlops</i> Heller, 1862
PENAEIDEA	Aristeidae	<i>Aristeomorpha foliacea</i> (Risso, 1827)
		<i>Aristeus antennatus</i> (Risso, 1816)
	Penaeidae	<i>Parapenaeus longirostris</i> (Lucas, 1846)
	Solenoceridae	<i>Solenocera membranacea</i> (Risso, 1816)
SERGESTOIDEA	Sergestidae	<i>Sergestes robustus</i> S.I.Smith, 1882
		<i>Sergestes articus</i> (Kröier, 1855)




Fig. 2. Occurrence and abundance indices of the collected crustaceans species (pooled surveys)

The occurrence values and biomass and density indices were determined for pooled data from the five surveys. The most prevalent species were the penaeid *Parapenaeus longirostris*, caridean shrimps of the genus *Plesionika*, portunid crabs *Liocarcinus depurator* and *Macropipus tuberculatus*, and astacidean *Nephrops norvegicus*, which accounted for most of total biomass in the area (Fig. 2) and were found at different depths (Fig. 3). *Liocarcinus depurator* was found on shelf bottoms while *P. longirostris*, *Plesionika* spp., and *M. tuberculatus* were found mostly on the shelf border and upper slope bottoms.

When carried out separately for the western and eastern areas, the analyses were slightly different. The western area was mostly characterized by the presence and abundance of Reptantia and burrowing species such as *L. depurator* and *N. norvegicus*, while the eastern area was characterized by Natantia species such as *P. longirostris* and *Plesionika* spp. (Figs. 4 and 5).

Sampling stations were clustered using a "species density per sampling station" matrix. Density indices per station were averaged for the overall period (all five surveys) to reduce the influence of occasional catches. Multivariate analysis resulted in four main clusters at the second level of ramification. The first cluster included most of the sampling stations deeper than 350 m (on both the western and eastern sides), the second one included mostly shelf and upper-slope stations in the eastern zone. The remaining two clusters included sampling stations in the western zone at 115-233 m and 21-125 m depths (Fig. 6).

According to the stratification of sampling scheme of the MEDITS protocol, classification and ordination (MDS) of data referring to the defined bathymetric strata (species mean density x depth strata, pooled surveys) per geographic area (southwestern and southeastern Adriatic) was performed. The differences between the western and eastern shelf areas were again revealed, while the slope zones appeared to be




Fig. 3. Bathymetric ranges and centroid distribution of the collected crustacean species




Fig. 4. Western zone: occurrence and abundance indices of the collected crustacean species (pooled surveys)




Fig. 5. Eastern zone: occurrence and abundance of indices the collected crustacean species (pooled surveys)




Fig. 6. Clustering of sampling stations from MEDITS 1996-2000 survey; crustaceans' density indices per station (no. km⁻²) were averaged on the overall period (five surveys); sampling stations are coded by mean depth and geographic area (W=western side; E=eastern side)




Fig. 7. Classification and ordination of crustacean assemblages on trawable bottoms of southern Adriatic Sea (W=western side; E=eastern side): mean density values per depth stratum (1=10-50 m; 2=51-100 m; 3=101-200 m; 4=201-500 m; 5=501-800 m); a) data averaged on the overall period (five surveys); b) data from 1996 MEDITS survey c) data from 2000 MEDITS survey

quite homogeneous (Fig. 7a). The same analysis was carried out for each yearly survey. Small inter-annual variations in the distribution of the main crustacean assemblages were detected only between the 1996 and 2000 surveys (Figs. 7b,c), confirming the persistence of assemblages in the investigated area.

The most abundant species on shelf bottoms in the western (*L. depurator* and *Goneplax rhomboides*) and eastern (*P. longirostris*) areas strongly contributed to the difference between the opposite sides of the southern Adriatic. The homogeneity on the slope bottoms was mostly due to the occurrence and abundance of *P. martia*, *Pasiphaea sivado*, *Aristaeomorpha foliacea*, and *Polycheles typhlops* on both sides.

DISCUSSION AND CONCLUSION

The list of species agrees with bibliographic information from the same geographic zone (PETRUZZI *et al.*, 1988; VASO & GJKNURI, 1993; PASTORELLI *et al.*, 1996; MARSAN *et al.*, 2000) and other Mediterranean areas, as well as the bathymetric ranges of distribution (RELINI-ORSI & RELINI, 1972; ABELLÓ *et al.*, 1988; RIGHINI & AUTERI, 1989; CARTES & SARDÀ, 1992; MURA & CAU, 1992; PIPITONE & TUMBILOLO, 1993; TURSI *et al.*, 1993; RINELLI *et al.*, 1998; SPANÒ, 1998). All the species in our study were included in lists regarding the whole Mediterranean basin (ŠTEVČIĆ & GALIL, 1994; FROGLIA, 1995). *Munida iris* Zariquiey Alvarez 1952 and *Calappa turkayana* PASTORE 1995 (PASTORE, 1995) were new findings for the southern Adriatic. They are added to *Paromola cuvieri* (RISSO, 1816) recently found in the same area (BELAMARIĆ *et al.*, 1995; UNGARO, 2000). The crustacean assemblages, as defined by multivariate analysis of the data, appeared to be rather stable over the years throughout the investigated area, as it is for most demersal species groups (GAERTNER *et al.*, 1998; UNGARO *et al.*, 1998).

The depth gradient can be very important for the distribution of fish assemblages in the southern Adriatic (UNGARO *et al.*, 1998, 1999) as it is worldwide but bathymetry, in addition, partially

explains the distribution in this area. Slope assemblages were homogeneously distributed throughout the area, while differences between the western and eastern zones were mostly on the shelf bottoms that are influenced by other variables as well as depth. The difference was probably due to geomorphic and oceanographic features, as the Adriatic Sea is a semi-enclosed basin characterized by latitudinal (south to north) as well as longitudinal (east to west) gradients (PIGORINI, 1968; ZORE-ARMANDA, 1968; ALFIREVIĆ, 1981; BREGANT *et al.*, 1992). Typical cyclonical movements are reported in the Adriatic (ARTEGIANI *et al.*, 1997) where dense cold water (north Adriatic dense water, NADW) flows from the north to the south along the western continental shelf, deep water (Adriatic deep water, ADW) forms in the southern Adriatic pit, and warmer more saline water (Levantine intermediate water, LIW) inflows northward along the eastern side of the Adriatic from the Ionian Sea in the Otranto Strait (MANCA *et al.*, 2001). The LIW makes the bottom water on the eastern side of the south Adriatic warmer than on the western side (ARTEGIANI *et al.*, 1997). Thus, the general water circulation in the basin as well as differences in the geomorphology can explain the distributional pattern of some of the crustacean species. On the other hand, a link between species distribution and talassographic parameters has been supposed (ABELLÓ *et al.*, 1988; CARTES *et al.*, 1994; UNGARO *et al.*, 1999), and new information from the southern Adriatic supports the probable role of bottom temperature for crustacean species distribution (UNGARO & MARANO, 2002; Fig. 8).

The results highlight the importance of research and study of the relationship between oceanographic features and the distribution of biological resources. Such study increases information on the marine ecosystem that can be used to manage the environment and exploitation of natural resources (e.g., fisheries).




Fig. 8. Relationship between abundance of *P. longirostris* and bottom temperature (mean values per survey, depths below 100 m; (W=western side; E=eastern side) (from UNGARO & MARANO, 2002)

REFERENCES

- ABELLÓ, P., F.J. VALLADARES & A. CASTELLÓN. 1988. Analysis of the structure of decapod crustacean assemblages off the Catalan coast (north-west Mediterranean). *Mar. Biol.*, 98: 39-49.
- ALFIREVIĆ, S. 1981. Contribution à la connaissance des caractéristiques bathymétriques et sédimentologiques de l'Adriatique. *FAO Fish. Rep.*, 253: 43-52.
- ARDISSON, P.L., E. BOURGET & P. LEGENDRE. 1990. Multivariate approach to study species assemblages at large spatiotemporal scales: The community structure of the epibenthic fauna of the Estuary and Gulf of St. Lawrence. *Can. J. Fish. Aquat. Sci.*, 47: 1364-1377.
- ARTEGIANI, A., D. BREGANT, E. PASCHINI, N. PINARDI, F. RAICICH & A. RUSSO. 1997. The Adriatic Sea general circulation. Part II: Baroclinic circulation structure. *J. Physic. Ocean.*, 27: 1515-1532.
- BELAMARIĆ, J.D., D. ZAVODNIK, E. ZAHTILA, A. NOVOSEL, N. OREPIĆ, I. OSOTIĆ, T. RADIŠA, M. RADOŠEVIĆ, J. ŠKALAMERA & J. VIDMAR. 1995. Preliminarni rezultati istraživanja makrobentosa uz vanjsku obalu Nacionalong parka "Mljet". *Ekološke monografije*, 6: 545-553.
- BERTRAND, J., L. GIL DE SOLA, C. PAPACOSTANTINO, G. RELINI & A. SOUPLET. 1997. An international bottom trawl survey in the Mediterranean: The MEDITS Programme. *ICES Ann. Sci. Conf. Y/03*, Baltimore C.M.
- BERTRAND, J.A., L. GIL DE SOLA, C. PAPACOSTANTINO, G. RELINI & A. SOUPLET. 2002. The general specifications of the MEDITS surveys. *Sci. Mar.*, 66(suppl. 2): 9-17.
- BREGANT, D., F. AZZARO, A. BONACCORSO, G. CIVITARESE, E. CRISAFI, R. LAFERLA, M. LEONARDI, A. LUCHETTA, R. POLIMENI & F. RACICH. 1992. Condizioni idrobiologiche nell'Adriatico Meridionale. *Atti X A.I.O.L.*, Alassio, Nov. 4-6: 37-46.
- CADDY, J.F. 1998. Issues in Mediterranean fisheries management: Geographical units and effort control. *Stud. Rev. G.F.C.M.*, 70: 1-56.
- CARR, M.R. 1996. *PRIMER User Manual (Plymouth Routines in Multivariate Ecological Research)*. Plymouth, U.K., 30 pp.
- CARTES, J.E. & F. SARDÀ. 1992. Abundance and diversity of decapod crustaceans in the deep Catalan sea (western Mediterranean). *Journal of Natural History*, 26: 1305-1323.
- CARTES, J.E., J.B. COMPANY & F. MAYNOU. 1994. Deep-water decapod crustacean communities

- in the north-western Mediterranean: Influence of submarine canyons and season, *Mar. Biol.*, 120: 221-229.
- CLARKE, K.R. & R.H. GREEN. 1988. Statistical design and analysis for a "biological effects" study. *Mar. Ecol. Prog. Ser.*, 46: 213-226.
- FALCIAI, L. & R. MINERVINI. 1992. Guida dei Crostacei Decapodi d'Europa. Franco Muzio (Editor), 282 pp.
- FROGLIA, C. 1995. Crustacea Malacostraca III (Decapoda). In: A. Minelli, S. Ruffo e S. La Posta (Editors). Checklist delle Specie della Fauna Italiana, Fascicolo 31, Edizioni Calderini Bologna.
- GAERTNER, J.C., D. CHESSEL & J. BERTRAND. 1998. Stability of spatial structures of demersal assemblages: a multitable approach, *Aquat. Living Resour.*, 11(2): 75-85.
- KRUSKAL, J.B. 1964. Multidimensional scaling by optimizing goodness of fit to a non-metric hypothesis. *Psychometrika*, 29: 1-27.
- LEGENDRE, L. & P. LEGENDRE. 1987. Developments in Numerical Ecology. NATO ASI Series G, Ecological Sciences 14, Springer-Verlag, Berlin, 585 pp.
- MANCA, B., P. FRANCO & E. PASCHINI. 2001. Seasonal variability of the hydrography in the Adriatic Sea: water mass properties and circulation. In: F.M. Faranda, L. Guglielmo, G. Spezie (Editors). Mediterranean Ecosystems Structures and Processes. Springer-Verlag, Milano, pp. 45-60.
- MARSAN, R., N. UNGARO, C.A. MARANO & M.C. MARZANO. 2000. Remarks on fishery biology of the genus *Plesionika* (Decapoda, Pandalidae) in the south Adriatic basin (Mediterranean Sea). *Crustacean Issues*, 12: 763-769.
- MURA, M. & A. CAU. 1992. Osservazioni su alcune comunità di vertebrati e macroinvertebrati demersali mesobatiali del canale di Sardegna. *Oebalia Suppl.*, XVII: 67-73.
- PASTORE, M. 1995. The genus *Calappa* in the Ionian Sea. *Oebalia*, XXI: 187-196.
- PASTORELLI, A.M., R. VACCARELLA, G. MARANO & N. UNGARO. 1996. I crostacei dei fondi strascicabili del Basso Adriatico. *Nova Thalassia*, 12: 27-35.
- PETRUZZI, T., A.M. PASTORELLI & G. MARANO. 1988. Notes on the distribution of commercial crustaceans in the southern Adriatic, Trawl-Survey 1985-86. *FAO Fish. Rep.*, 394: 213-221.
- PIGORINI, B. 1968. Sources and dispersion of recent sediments of the Adriatic Sea. *Mar. Geol.*, 6: 187-229.
- PIPTONE, C. & M.L. TUMBILOLO. 1993. Decapod and stomatopod crustaceans from the trawlable bottoms of the Sicilian Channel (central Mediterranean Sea). *Crustaceana*, 65(3): 358-364.
- ORSI, L. & G. RELINI. 1972. Note sui crostacei decapodi batiali del Mar Ligure. *Boll. Mus. Ist. Biol. Univ. Genova*, 40: 47-73.
- RELINI, G., J. BERTRAND & A. ZAMBONI. 1999. Synthesis of the knowledge on bottom fishery resources in central Mediterranean (Italy and Corsica). *Biol. Mar. Medit.*, 6(suppl. 1): 868 pp.
- RIGHINI, P. & R. AUTERI. 1989. Distribuzione batimetrica dei crostacei decapodi raccolti durante le campagne di pesca nel Tirreno meridionale. *Oebalia N.S.*, 15(2): 763-767.
- RINELLI, P., N. SPANÒ & S. GRECO. 1998. Distribuzione dei crostacei decapodi ed echinodermi dei fondi strascicabili del Mar Tirreno meridionale. *Biol. Mar. Medit.*, 5(2): 211-217.
- SPANÒ, N. 1998. Decapoda crustacea in the southern Tyrrhenian Sea – five years of research. *Rapp. Comm. int. Mer Médit.*, 35: p. 488.
- SPARRE, P. & S.C. VENEMA. 1992. Introduction to tropical fish stock assessment. *FAO Fish. Tech. Pap.*, 306/1(rev. 1): 1-376.
- ŠTEVČIĆ, Z. & B. GALIL. 1994. Checklist of the Mediterranean brachyuran crabs. *Acta Adriat.*, 34(1/2): 65-76.
- TURSI A., G. DONGHIA, A. MATARRESE, C. CAROPPO & G. COSTANTINO. 1993. L'importanza dei crostacei decapodi (natanti e reptanti macruri) nel contesto delle campagne di pesca condotte nel Mar Ionio (1985-1986). *Quaderni dell'Istituto Ricerche Pesca Marittima*, 5(2): 145-158.
- UNGARO, N. 2000. Ritrovamenti di *Paromola cuvieri* (Risso, 1816) (Dromopoda, Homolidae) nel bacino Adriatico meridionale. *Biol. Mar. Medit.*, 7(2): 744-747.

- UNGARO, N. & G. MARANO. 2001. Fish species biodiversity on trawlable bottoms of south Adriatic basin (Mediterranean Sea): years 1996-1998. In: F.M. Faranda, L. Guglielmo, G. Spezie (Editors). Structures and Processes in the Mediterranean Ecosystems. Springer-Verlag, Milano, pp. 215-220.
- UNGARO, N. & G. MARANO. 2002. On the distribution and demography of the deep-water pink shrimp and Norway lobster fishery stocks in the south Adriatic Sea. *Studia Marina*, 23(1): 15-24.
- UNGARO N., G. MARANO & R. VACCARELLA. 1995. Comparazione tra aree batiali strascicabili del basso Adriatico mediante l'utilizzo dell'analisi fattoriale delle corrispondenze. *Biol. Mar. Med.*, 2(2): 185-189.
- UNGARO, N., G. MARANO, A. VLORA & M. MARTINO. 1998. Space-time variations of demersal fish assemblages in south-western Adriatic Sea. *Vie Milieu*, 48(3): 191-201.
- UNGARO, N., C.A. MARANO, R. MARSAN, M. MARTINO, M. MARZANO, G. STRIPPOLI & A. VLORA. 1999. Analysis of demersal species assemblages from trawl surveys in the south Adriatic Sea. *Aquat. Living Resour.*, 12(3): 177-185.
- VASO, A. & L. GJIKNURI. 1993. Decapod crustaceans of the Albanian coast. *Crustaceana*, 65(3): 390-407.
- ZARIQUIEY-ALVAREZ, R. 1968. Crustáceos decápodos Ibéricos. *Inv. Pesqu.*, 32: 1-510.
- ZORE-ARMANDA, M. 1968. The system of currents in the Adriatic Sea. *Rev. Fish. Counc. Medit.*, 34: 1-48.

Received: 13 January 2004

Accepted: 17 September 2004

Raspodjela demerzalnih rakova u južnom Jadranu

Nicola UNGARO ¹, Chiara Alessandra MARANO ², Luca CERIOLA ¹ i Michele MARTINO ²

¹*Pokrajinski laboratorij za biologiju mora, Molo Pizzoli (Porto), 70123 Bari, Italija*

²*Odjel za životinjsku proizvodnju, Poljoprivredni fakultet Sveučilišta, Via Amendola 126/a, 70126 Bari, Italija*

SAŽETAK

Iz podataka sabranih tijekom pet sezonskih krstarenja (ljeti) koćom od 1996 do 2000. godine, određeni su pojava i raspodjela rakova, te njihove zajednice u vodama južnog Jadrana. Koćarenje je bilo izvedeno u okviru E.U. projekta MEDITS, na dubinama od 10-800 m, istim brodom i alatom. Ulovljene su 52 bentoske-epibentoske i nektonske vrste (3 stomatopoda, 49 dekapoda). Podaci o gustoći vrsta (primjeraka na km²) obrađeni su multivarijantnim tehnikama kako bi se utvrdio sastav i raspodjela glavnih zajednica rakova. Zajednice su bile jednake u čitavom istraživanom području, s nekim razlikama između zapadnih i istočnih zona, najčešće na području šelfa. Diskutiran je i utjecaj oceanografskih svojstava osim dubine.

Ključne riječi: dekapodni raci, stomatopodi, demerzalne zajednice, multivarijantna analiza, Jadransko more