

Original Scientific Paper

## SUBSTITUTED MONOQUATERNARY OXIMES AS REACTIVATORS OF CYCLOSARIN- AND CHLORPYRIFOS-INHIBITED ACETYLCHOLINESTERASE

Veronika RACAKOVA<sup>1</sup>, Martina HRABINOVA<sup>2</sup>, Daniel JUN<sup>2</sup> and Kamil KUČA<sup>2</sup>

*Department of Pharmaceutical Chemistry and Drug Control, Faculty of Pharmacy, Charles University of Prague<sup>1</sup>, Department of Toxicology, Faculty of Military Health Sciences, University of Defence<sup>2</sup>, Hradec Kralove, Czech Republic*

Received in March 2006

Accepted in August 2006

This paper describes an *in vitro* study of three potential acetylcholinesterase (AChE; EC 3.1.1.7) reactivators derived from a monoquaternary reactivator pralidoxime. Compounds used were pyridinium-2-aldoxime-4-carbamoyl-N-methyl iodide (TO231), pyridinium-2-aldoxime-4-ethoxycarbonyl-N-methyl iodide (TO237), and pyridinium-2-aldoxime-5-ethoxycarbonyl-N-methyl iodide (TO238). Pralidoxime and obidoxime were used for comparison. Nerve agent cyclosarin and pesticide chlorpyrifos were used as organophosphorus cholinesterase inhibitors. The source of AChE was rat brain homogenate. None of the tested oximes was able to reactivate cyclosarin-inhibited AChE (at 1.0 mmol L<sup>-1</sup> oxime concentration). In case of chlorpyrifos, TO231 was the most potent AChE reactivator with an 82 % reactivation at 1.0 mmol L<sup>-1</sup> oxime concentration. This reactivating potency equals that of pralidoxime and obidoxime. TO238 was less effective, and TO237 did not reactivate chlorpyrifos-inhibited AChE at all. None of the tested AChE reactivators, reference compounds included, could be considered universal for both chlorpyrifos- and cyclosarin-inhibited AChE.

**KEY WORDS:** *brain cholinesterase, cholinesterase inhibition, nerve agents, organophosphorus compounds, pesticides, poisoning, reactivation, treatment*

The toxicity of organophosphorus (OP) compounds - which are used in agriculture as pesticides, in medicine as drugs and in the military as nerve agents - is mostly related to the inhibition of acetylcholinesterase (AChE; EC 3.1.1.7), a very important enzyme which splits the neuromediator acetylcholine (ACh) at the synaptic clefts. Generally, anticholinergics (as functional drugs) and AChE reactivators (as causal drugs) are used as the first aid antidotes in OP poisoning (1-3).

In contrast to anticholinergics that can be administered in case of any nerve agent poisoning, a need has arisen for specific AChE reactivators to counter AChE inhibition. Although there are many potential AChE reactivators, none is able to reactivate

all OP inhibitions, regardless of the inhibitor structure (4-6). This is why many laboratories throughout the world are interested in finding a new, broad-spectrum AChE reactivator (7-11). Currently the most promising AChE reactivator HI-6 has a limited spectrum due to its low efficacy in reactivating tabun- and pesticide-inhibited AChE (12, 13). Pralidoxime, a gold standard among AChE reactivators (14), which is presently used as antidote in the US, is a very poor reactivator of AChE inhibited by a nerve agent (15-17).

This study evaluated monoquaternary oximes TO238, TO231, and TO237, prepared in our department (18). Although these oximes are known for more than eight years, they have never been tested

as AChE reactivators after cyclosarin or chlorpyrifos poisoning. We studied these compounds as *in vitro* reactivators of cyclosarin- and chlorpyrifos-inhibited AChE using our general potentiometric method (19).

## MATERIAL AND METHODS

All tested compounds were prepared earlier at the Department of Toxicology of the Faculty of Military Health Sciences (Hradec Kralove; Czech Republic) (18). Their structures are listed in Figure 1. Purities of the tested AChE reactivators were established using TLC (DC-Alufolien Cellulose F; Merck, Germany; mobile phase BuOH-CH<sub>3</sub>COOH-H<sub>2</sub>O 5 : 1 : 2; detection by solution of Dragendorff reagent). Chlorpyrifos [O,O-diethyl-O-(3,5,6-trichloro-2-pyridyl)-phosphorothioate] was obtained from Riedel-de Haen as analytical standard (99.2 % purity). Nerve agent cyclosarin (GF; O-cyclohexylmethylflourophosphate) was obtained from the Military Facility Brno (97 % purity). All other chemicals used were of reagent grade (Sigma-Aldrich, Czech Republic).

Rat brain homogenate was incubated with the OP compound for 30 min and the AChE reactivator was then added. Ten minutes later the substrate (ACh) was added and the activity measurement started. The released acetic acid from ACh was titrated using sodium hydroxide. The consumption of sodium hydroxide was proportional to the enzyme activity, allowing us to calculate the reactivation potency of the tested AChE reactivators. The whole *in vitro* experiment (including the equation used to calculate

the percent of reactivation) was described in detail earlier (19, 20).

## RESULTS AND DISCUSSION

Table 1 shows that AChE reactivators TO231 and TO238 had a higher reactivation potency against the pesticide chlorpyrifos than against cyclosarin. Only obidoxime and TO231 achieved a low reactivation potency against cyclosarin (below 10 %). For chlorpyrifos-inhibited AChE, pralidoxime, obidoxime, TO231, and TO238 reached a promising reactivation potency at the concentration of 1.0 mmol L<sup>-1</sup>. Only obidoxime showed a promising reactivation potency at the concentration of 10.0 μmol L<sup>-1</sup> and was thus the most promising oxime in this study.

There is no single AChE reactivator able to serve as a broad-spectrum antidote against both nerve agent and pesticide poisoning. Currently the most promising oxime HI-6 does not sufficiently reactivate tabun- and pesticide-inhibited AChE (12, 13). This is why attempts to synthesize more effective oximes continue (7-11). To find an effective AChE reactivator new compounds are being developed (21, 22). However none of these substances, tested in a number of *in vitro* and *in vivo* studies, can be considered broad-spectrum reactivators (12, 13, 23, 24).

We therefore focused our attention on AChE reactivators that had been known for more than eight years, but had been never tested for their broad-spectrum reactivation potency. Owing to the fact that the only experiments done with these oximes were conducted on mice using a pesticide fosdrine,




Figure 1 Structure of tested AChE reactivators

**Table 1** Reactivation potency of oximes tested (enzyme source: rat brain homogenate; time of inhibition: 30 min; time of reactivation: 10 min; pH 7.6; 25 °C; n=3)

Oxime	ORGANOPHOSPHORUS COMPOUND			
	Chlorpyrifos		Cyclosarin	
	Reactivation potency / % *	Reactivation potency / % #	Reactivation potency / % *	Reactivation potency / % #
Pralidoxime	9	80	0	0
Obidoxime	43	76	5	2
TO238	0	48	0	0
TO231	6	82	0	3
TO237	0	0	0	0

\* oxime concentration = 10.0 μmol L<sup>-1</sup>

# oxime concentration = 1.0 mmol L<sup>-1</sup>

we decided to test them against the nerve agent cyclosarin and a pesticide group member chlorpyrifos (18). Our results showed that none of the tested oximes was able to reactivate cyclosarin-inhibited AChE, but some oximes reached promising reactivation potency against chlorpyrifos at the concentration of 1 mmol L<sup>-1</sup>. Our results suggest that there is no point in investigating these reactivators further in order to obtain a broad-spectrum reactivator.

#### Acknowledgement

This study was supported by a Charles University (Czech Republic) grant No. 126/2005/B-BIO/FaF and by a Ministry of Defence (Czech Republic) grant No. FVZ0000501.

#### REFERENCES

- Bajgar J. Organophosphates/nerve agent poisoning: mechanism of action, diagnosis, prophylaxis, and treatment. *Adv Clin Chem* 2004;38:151-216.
- Patocka J, Kuca K, Jun D. Acetylcholinesterase: crucial enzyme of human body. *Acta Medica (Hradec Kralove)* 2004;47:215-30.
- Marrs TC. Organophosphate poisoning. *Pharmacol Therap* 1993;58:51-66.
- Sevelova L, Vachek J. Effect of methoxime combined with anticholinergic, anticonvulsant or anti-HCN drugs in tabun-poisoned mice. *Acta Medica (Hradec Kralove)* 2003;46:109-12.
- Shih TM, Koviak TA, Capacio BR. Anticonvulsants for poisoning by the organophosphorus compound soman: pharmacological mechanisms. *Neurosci Biobehav Rev* 1991;15:349-62.
- Kassa J. Review of oximes in the antidotal treatment of poisoning by organophosphorus nerve agents. *J Toxicol Clin Toxicol* 2002;40:803-16.7.
- Picha J, Kuca K, Kivala M, Kohout M, Cabal J, Liska F. New group of monoquaternary reactivators of the acetylcholinesterase inhibited by nerve agents. *J Enzyme Inhib Med Chem* 2005;20:233-7.
- Musilek K, Kuca K, Jun D, Dohnal V, Dolezal M. Synthesis of the novel series of bispyridinium compounds bearing xylene linker and evaluation of their reactivation activity against chlorpyrifos-inhibited acetylcholinesterase. *J Enzyme Inhib Med Chem* 2005;20:409-15.
- Kim TH, Kuca K, Jun D, Jung YS. Design and synthesis of new bis-pyridinium oximes as cyclosarin-inhibited acetylcholinesterase reactivators. *Bioorg Med Chem Lett* 2005;15:2914-7.
- Yang GY, Yoon JH, Seong CM, Park NS, Jung YS. Synthesis of bis-pyridinium oxime antidotes using bis(methylsulfonylmethyl) ether for organophosphate nerve agents. *Bull Korean Chem Soc* 2003;24:1368-70.
- Chennamaneni SR, Vobalaboina V, Garlapati A. Quaternary salts of 4,3' and 4,4' bis-pyridinium monooximes: Synthesis and biological activity. *Bioorg Med Chem Lett* 2005;15:3076-80.
- Calic M, Lucic Vrdoljak A, Radic B, Jelic D, Jun D, Kuca K, Kovarik Z. In vitro and in vivo evaluation of pyridinium oximes: mode of interaction with acetylcholinesterase, effect on tabun- and soman-poisoned mice and their cytotoxicity. *Toxicology* 2006;219:85-96.
- Worek F, Kirchner T, Backer M, Szinicz L. Reactivation by various oximes of human erythrocyte acetylcholinesterase inhibited by different organophosphorus compounds. *Arch Toxicol* 1996;70:497-503.
- Petroianu GA, Hasan MY, Nurulain SM, Arafat K, Shafiullah M, Naseer O. Protective agents in acute high-dose organophosphate exposure: comparison of ranitidine with pralidoxime in rats. *J Appl Toxicol* 2005;25:68-73.
- Kuca K, Cabal J, Kassa J. A comparison of the efficacy of a bispyridinium oxime – 1,4-bis-(2-

- hydroxyiminomethylpyridinium) butane dibromide and currently used oximes to reactivate sarin, tabun or cyclosarin-inhibited acetylcholinesterase by *in vitro* methods. *Die Pharmazie* 2004;59:795-8.
16. Worek F, Diepold C, Eyer P. Dimethylphosphoryl-inhibited human cholinesterases: inhibition, reactivation, and aging kinetics. *Arch Toxicol* 1999;73:7-14.
  17. Sevelova L, Kuca K, Krejčova G. Antidotal treatment of GF-agent intoxication in mice with new bispyridinium oximes. *Toxicology* 2005;207:1-6.
  18. Bielavsky J, Kassa J, Elsnerova I, Dejmek L. Cholinesterase reactivators derived from pyridine-2-carbaldoxime. *Coll Czech Chem Commun* 1998;63:199-204.
  19. Kuca K, Cabal J. Evaluation of newly synthesized reactivators of the brain cholinesterase inhibited by sarin-nerve agent. *Toxicol Mech Meth* 2005;15:247-52.
  20. Kuca K, Kassa J. A Comparison of the ability of a new bispyridinium oxime-1-(4-hydroxyiminomethylpyridin-ium)-4-(4-carbamoylpyridinium)butane dibromide and currently used oximes to reactivate nerve agent-inhibited rat brain acetylcholinesterase by *in vitro* methods. *J Enzyme Inhib Med Chem* 2003;18:529-35.
  21. Eyer P, Ladstetter B, Schafer W, Sonnenbichler J. Studies on the stability and decomposition of the Hagedorn-oxime HLo 7 in aqueous solution. *Arch Toxicol* 1989;63:59-67.
  22. Kuca K, Bielavsky J, Cabal J, Bielavska M. Synthesis of a potential reactivator of acetylcholinesterase 1-(4-hydroxyiminomethylpyridinium)-3-(carbamoylpyridinium)-propane dibromide. *Tetrahedron Lett* 2003;44:3123-5.
  23. Eyer P. The role of oximes in the management of organophosphorus pesticide poisoning. *Toxicol Rev* 2003;22:165-90.
  24. Kuca K, Kassa J. In vitro reactivation of acetylcholinesterase using of the oxime K027. *Vet Hum Toxicol* 2004;46:15-8.

### Sažetak

#### SUPSTITUIRANI MONOKVATERNI OKSIMI KAO REAKTIVATORI ACETILKOLINESTERAZE INHIBIRANE CIKLOSARINOM I KLORPIRIFOSOM

U ovome *in vitro* istraživanju testiran je reaktivacijski potencijal triju reaktivatora acetilkolinesteraze (AChE; EC 3.1.1.7) dobivenih iz monokvaternog reaktivatora pralidoksima. Testirani su sljedeći spojevi: 4-karbamoil-2-aldoksim-1-metilpiridinijev jodid (TO231), 4-etoksikarbonil-2-aldoksim-1-metilpiridinijev jodid (TO237) te 5-etoksikarbonil-2-aldoksim-1-metilpiridinijev jodid (TO238). Za usporedbu su uzeti pralidoksim i obidoksim. Kao inhibitori kolinesteraze rabljeni su živčani otrov ciklosarin i pesticid klorpirifos. Izvor acetilkolinesteraze bio je homogenat mozga štakora. Nijedan od testiranih oksima nije uspio reaktivirati acetilkolinesterazu inhibiranu ciklosarinom pri koncentraciji oksima od 1.0 mmol L<sup>-1</sup>. Kod inhibicije klorpirifosom TO231 pokazao se najsnažnijim reaktivatorom acetilkolinesteraze s reaktivacijom od 82 % pri koncentraciji oksima od 1.0 mmol L<sup>-1</sup>. Ovaj reaktivacijski potencijal jednak je onomu pralidoksima i obidoksima. Slabiju je djelotvornost iskazao TO238, a TO237 nije uspio reaktivirati acetilkolinesterazu inhibiranu klorpirifosom. Stoga nijedan od testiranih spojeva (uključujući i referentne oksime) ne može služiti kao univerzalni reaktivator acetilkolinesteraze inhibirane klorpirifosom i ciklosarinom.

**KLJUČNE RIJEČI:** *inhibicija kolinesteraza, kolinesteraza mozga, liječenje, organofosforni spojevi, pesticidi, živčani otrovi*

#### REQUESTS FOR REPRINTS:

Kamil KUČA  
Department of Toxicology  
Faculty of Military Health Sciences  
University of Defence  
Trebesska 1575, 500 01, Hradec Kralove, Czech Republic  
E-mail: [kucakam@pmfhk.cz](mailto:kucakam@pmfhk.cz)