

THE INFLUENCE OF THE FLOATER POSITION ON THE LOAD OF SOCCER PLAYERS DURING A 4 VS 4 + 2 GAME

Jose Antonio Asian-Clemente ^{1,2,3}, Alberto Rabano-Muñoz ³, Bernardo Requena ¹, Alfredo Santalla ^{1,2}, and Luis Suarez-Arrones ^{1,2}

¹Football Science Institute, Granada, Spain

²Department of Sport Sciences, University Pablo de Olavide, Sevilla, Spain

³Real Betis Balompié, Sevilla, Spain

Original scientific paper

DOI 10.26582/k.54.1.9

Abstract:

The purpose of this study was to examine the load of regular players and floater players in a specific small-sided game. Twenty semi-professional soccer players performed one 4vs4+2, modifying the position of the floaters (internal, external, zone, square and without floaters). Total distance covered, distance covered at speeds between 7-13.9 km·h⁻¹, 14-17.9 km·h⁻¹, and ≥18 km·h⁻¹, accelerations and decelerations between 2.5-4 m·s⁻², maximal and mean heart rate (HR) and rating of perceived exertion (RPE) were analyzed. Internal floaters achieved greater total distance covered, accelerations and RPE than in any other position. Internal and external floaters achieved more distance covered at 7-13.9, 14-17.9, >18 km·h⁻¹ and HR than zone and square floaters. With internal floaters, regular players covered more distance covered >18 km·h⁻¹ than in any other 4vs4+2 format, and with internal and external floaters regular players covered greater total distance and distance at 14-17.9 km·h⁻¹ than without floaters or with zone or square floaters. Regular players showed greater total distance covered, distance covered at 7-13.9, 14-17.9, >18 km·h⁻¹, accelerations and RPE than floaters in all 4vs4+2. These data showed that floater position in 4vs4+2 game influenced the internal and external load of both the regular players and the floaters.

Key words: football, time-motion, global position system (GPS), heart rate, rate of perceived exertion

Introduction

Soccer is a sport that can be considered as a set of complex, self-organized, unstable, unpredictable and highly dynamic systems in which players interact with each other trying to achieve offensive and defensive maneuvers (Davids, Araujo, Correia, & Vilar, 2013; Vilar, Araujo, Davids, & Button, 2012). Sometimes, that interaction can be understood as random and stochastic (Low, et al., 2020; Mackenzie & Cushion, 2013; Yue, Broich, Seifriz, & Mester, 2008), so one of the most important issues in enhancing performance in soccer is the incorporation of training drills that totally or partially mimic the requirements of a formal competition (Katis & Kellis, 2009), to obtain stimuli that would provide the maximum benefits for the soccer player with regard to real-world matches (Mallo & Navarro, 2008; Turner & Stewart, 2014).

Small sided games (SSGs) are among the most frequently applied and most popular training exercises used by soccer coaches (Clemente, et al., 2020; Clemente, Praca, Bredt, van der Linden, & Serra-Olivares, 2019; Owen, Newton, Shovlin, &

Malone, 2020; Praca, Andrade, Bredt, Moura, & Moreira, 2022; Rampinini, et al., 2007; Sarmiento, et al., 2018) because, like matches, SSGs provide players with opportunities for continuous exploration, discovery and learning of specific situations (Ric, Hristovski, & Torrents, 2015; Rico-Gonzalez, Pino-Ortega, Praca, & Clemente, 2022) while simultaneously aiming at the development of technical, tactical, and physical components (Dellal, et al., 2008; Hill-Haas, Rowsell, Dawson, & Coutts, 2009). SSGs are modified games played on a reduced-size field, often involving a smaller number of players and introducing adapted rules, which are different from regular eleven-a-side games (Hill-Haas, Dawson, Impellizzeri, & Coutts, 2011). In recent years, the available research on manipulating the game rules applied during SSGs has increased significantly (Aguilar, Botelho, Lago, Macas, & Sampaio, 2012; Castillo, Raya-Gonzalez, Manuel Clemente, & Yanci, 2020; Coutinho, et al., 2020; Hill-Haas, et al., 2011; Sarmiento, et al., 2018). Most studies have been carried out in teams with the same numbers of players (Torres-Ronda, et al.,

2015) even though soccer is frequently played in numerical imbalance, either temporarily or permanently (Sampaio, Lago, Goncalves, Macas, & Leite, 2014), and despite the fact that coaches regularly use floater players to regulate the load on some players (Lacome, Simpson, Cholley, & Buchheit, 2018). Only a few studies have investigated the internal and external loads during SSGs of such formats. Hill-Haas, Coutts, Dawson, and Rowsell (2010) showed that floaters covered greater total distances and engaged in more sprints than regular players (RP), comparing various SSG formats with different number of players, dissimilar pitch sizes and rule changes. Only two of four SSG formats used had one floater, so we cannot compare floaters' vs RPs' demands in the results obtained in Hill-Haas et al.'s study (Hill-Haas, et al., 2010). Lacome et al. (2018) recently demonstrated that locomotor activity and external mechanical load were lower in floaters than in RP in SSGs with one floater, depending on pitch size variations, which modified the number of players, and the inclusion of a goalkeeper. Keeping the total size of the field but modifying the length-width orientation in SSGs, Gollin, Alfero, and Abate Daga (2016) detected a lower load in internal floaters than in RP. Similar results have been found for internal floaters when the orientation of the SSGs is kept (Rabano-Munoz, Asian-Clemente, Saez de Villarreal, Nayler, & Requena, 2019). Internal load parameters measured as rating of perceived exertion (RPE) and heart rate (HR), and technical demands have been also analyzed, comparing SSG + 2 internal floaters, SSG + 2 internal and 2 external floaters, and SSG without floaters (Sanchez-Sanchez, et al., 2017). However, we are not aware of any study that has compared floaters' internal and external loads with those of RPs depending on their position in SSG formats with the same pitch size, the number of players and rules. Since it is common for coaches to change floaters' positions during SSGs, it is of interest to determine the effect of the floater role on RP and floater's demands during SSGs, isolating this from other influencing variables.

Therefore, the purpose of this study was to examine acute physiological responses and time-motion characteristics of RPs and floaters, and their relationship to the position of the floater, during a 4vs4+2 with the same pitch size, the number of players and rules.

Material and methods

Participants

Twenty semi-professional male soccer players (age: 24.1 ± 3.5 years; body height: 177.2 ± 5.9 cm; body weight: 70.3 ± 8.2 kg; % body fat (Faulkner, 1968): 11.5 ± 1.9) from the same team participated

in the study. All players were informed on the aims of the study and gave their informed consent before starting. The study was approved by the local ethical committee of the University of Pablo de Olavide and was developed according to the principles of the Declaration of Helsinki for the study with human subjects.

Procedures

Data were collected during the first half of the competitive season 2017-2018. Players were involved in five training sessions per week (80-120 minutes) plus one competitive match (played on Sunday). To guarantee the recovery of the players, measurements took place at the same time of the week after the rest day (Wednesday). All these sessions started by assigning the same GPS to the players to limit measurement errors, continued with a similar 20-min standardized warm-up and were completed on the natural turf at the same time of a day. After the warm-up, the players' behavior in SSGs was monitored and the session finished with a conditioned game with high technical and tactical requirements.


Small-sided games

Five SSGs were implemented over the duration of the study (Figure 1). On each drill monitored a representative moment of the game was represented. Training tasks were designed by technical staff to ensure the transference to the match by following specificity principle. The teams in the SSGs were balanced according to players' technical and tactical level, competitive experience, playing positions (Dellal, et al., 2012) and the subjective evaluation of the coaches (Casamichana & Castellano, 2010). The role of floaters was assigned to the same players during the study, and it was chosen by the coaches.

Teams of four players aimed to retain ball possession, with the incorporation of two floaters, who always assumed an offensive role (Campos-Vazquez, et al., 2017). The pitch size (40 x 30 m) and the duration (four bouts of 4-min, separated by 2-min of passive recovery) were constant in all the SSGs (Kelly & Drust, 2009; Sanchez-Sanchez, et al., 2017). SSGs were performed with no limit on ball touches and no player-to-player marking. Coaches verbally encouraged the players to maintain a high work rate during the SSG bouts.

Locomotor activity

Running activity parameters were monitored using a GPS system (GPSports SPI Elite System, Canberra, Australia) previously validated in the literature (Barbero-Alvarez, Coutts, Granda, Barbero-Alvarez, & Castagna, 2010; Coutts & Duffield, 2010). Total distance covered (TD), and



SSGs format: a) 4vs4; b) 4vs4 + 2 internal floaters; c) 4vs4 + 2 external floaters; d) 4vs4 + 2 zone floaters; e) 4vs4 + 2 square floaters. **Grey points: regular players in possession of the ball. Black triangles: regular players defending.** White points: floaters (attacking). RARP = Relative area regular players; RAFF = Relative area floater players

Figure 1. Graphical representation of the formats of the SSGs.

distances covered (DC) at several speeds: DC at 7-13.9 km·h⁻¹, 14-17.9 km·h⁻¹, and ≥ 18 km·h⁻¹ (Casamichana, Castellano, & Castagna, 2012). Accelerations and decelerations between 2.5-4 m·s⁻² were also recorded (Buchheit, et al., 2014; Suarez-Arrones, et al., 2016).

Heart rate (HR) and rate of perceived exertion (RPE)

Maximal (HR_{max}) and mean (HR_{mean}) heart rates achieved by players during the SSGs were obtained using a short-range radio telemetry (1 Hz, Polar Team Sports System, Polar Electro Oy, Finland) and integrated in the GPS units across the games. The 1-10 Borg Scale for RPE was used to assess the global SSG internal load (Borg, Hassmen, & Lagerstrom, 1987; Foster, et al., 2001). Players, having already been familiarized with that scale, gave their evaluations immediately after each game using a standardized questionnaire (Coutts, Rampinini, Marcora, Castagna, & Impellizzeri, 2009).

Statistical analyses

Data are presented as mean ± standard deviation (SD). All variables showed a normal distribution (Shapiro-Wilk test). A one-way analysis of variance (ANOVA) was used to determine variations in the TD, DC in each speed zone, number of accelerations and decelerations, HR and RPE. Cohen's effect size (ES) was also calculated to compare the magnitudes of the change between groups in certain variables, and quantitative differences were assessed

(Hopkins, Marshall, Batterham, & Hanin, 2009). Threshold values for assessing magnitudes of the ES were < 0.20, 0.20, 0.60, 1.2, and 2.0 for trivial, small, moderate, large, and very large, respectively (Hopkins, et al., 2009). The SPSS statistical software package (V20.0 for Windows, SPSS Inc., Chicago, IL) was used for data analysis.


Results

Comparisons between floater players

Table 1 and Figure 2 show the external and internal loads of floaters depending on their position in the 4vs4+2 SSGs. Total distance (TD), number of accelerations and RPE were statistically higher in the internal floaters than in any other position (very large to small magnitude). Likewise, internal floaters achieved a statistically greater number of decelerations than the external and square floaters; and greater DC at 7-13.9, 14-17.9, > 18 km h⁻¹, HR_{mean} and HR_{max} than the zone and square floaters (large to small magnitude). External floaters performed statistically more DC at 7-13.9, 14-17.9, > 18 km h⁻¹, HR_{mean} and RPE than the zone and square floaters (large to small magnitude), more TD, accelerations and decelerations than the square floaters (large and moderate magnitude) and fewer decelerations than the zone floaters (small magnitude). There were also statistical differences between the zone and square floaters, with the zone floaters showing more TD, DC at 7-13.9 and 14-17.9 km·h⁻¹, accelerations and decelerations (large to moderate magnitude).

Table 1. Descriptive data for external and internal load of the floaters in the 4vs4+2.

Variable	Internal Floaters	External floaters	Zone Floaters	Square Floaters
TD (m)	1847.4 ± 120.3	1012.8 ± 14.9	960.7 ± 207.6	763.4 ± 93.5
DC 7–13.9 km·h ⁻¹	431.5 ± 284.8	275.0 ± 25.0	157.4 ± 108.4	74.4 ± 42.2
DC 14–17.9 km·h ⁻¹	65.1 ± 12.5	25.2 ± 15.0	5.9 ± 6.8	1.4 ± 0.35
DC ≥ 18 km·h ⁻¹	8.8 ± 6.4	3.4 ± 3.9	0.0 ± 0.0	0.0 ± 0.0
Acc 2.5–4 m·s ⁻²	13.0 ± 3.7	8.0 ± 1.2	8.5 ± 1.7	2.0 ± 1.15
Dec 2.5–4 m·s ⁻²	11.3 ± 4.9	5.1 ± 1.4	8.0 ± 2.1	2.2 ± 1.4
HR _{mean}	147.9 ± 18.6	147.3 ± 17.0	138.6 ± 6.9	136.4 ± 18.6
HR _{max}	180.0 ± 12.4	172.5 ± 17.7	170.0 ± 8.3	170.5 ± 12.0
RPE	6.8 ± 0.3	5.8 ± 0.4	4.8 ± 0.4	4.75 ± 0.4




Bars indicate uncertainty in the true mean changes with 90% confidence intervals. Trivial (shaded) areas were calculated from the smallest worthwhile change. TD = Total distance covered; DC = Distance covered; Acc = accelerations; Dec = decelerations; HR = heart rate; RPE = rate of perceived exertion; ES = Effect size; * = p < 0.05; ** p < 0.01

Figure 2. Comparison of the external and internal load of floater players depending on their positions in the SSG

Comparisons between regular players (RP)

Table 2 and Figure 3 present the internal and external load of RPs for the proposed SSGs. With

internal floaters, RPs covered statistically more DC at > 18 km·h⁻¹ than in any other SSG format (very large and large magnitude). With internal and external floaters, RPs reached statistically



Bars indicate uncertainty in the true mean changes with 90% confidence intervals. Trivial (shaded) areas were calculated from the smallest worthwhile change. TD = Total distance covered; DC = Distance covered; Acc = accelerations; Dec = decelerations; HR = heart rate; RPE = rate of perceived exertion; ES = Effect size; * = p < 0.05; ** p < 0.01

Figure 3. Comparison of external and internal load of RP in different SSGs.

Table 2. Descriptive data for external and internal load of RP in SSGs.


Variable	4vs4		4vs4+2		
	Without floaters	Internal Floaters	External floaters	Zone Floaters	Square Floaters
TD (m)	1826.4 ± 107.2	1975.2 ± 169.3	1982.0 ± 135.4	1862.7 ± 191.8	1859.6 ± 155.0
DC 7–13.9 km·h ⁻¹	857.3 ± 90.4	943.6 ± 160.3	1005.6 ± 163.0	765.7 ± 306.0	915.6 ± 155.3
DC 14–17.9 km·h ⁻¹	213.2 ± 73.6	293.7 ± 90.5	271.7 ± 54.8	219.6 ± 91.0	216.8 ± 70.6
DC ≥ 18 km·h ⁻¹	16.2 ± 10.7	46.6 ± 19.6	20.7 ± 16.5	18.1 ± 14.7	16.4 ± 12.2
Acc 2.5–4 m·s ⁻²	21.6 ± 4.4	18.0 ± 3.4	21.3 ± 7.2	16.4 ± 2.6	19.1 ± 5.9
Dec 2.5–4 m·s ⁻²	15.7 ± 3.6	13.3 ± 4.4	19.8 ± 8.4	14.9 ± 3.8	24.0 ± 5.9
HR _{mean}	150.5 ± 44.3	151.2 ± 16.4	159.7 ± 15.6	156.3 ± 17.2	150.8 ± 18.8
HR _{max}	177.4 ± 21.3	181.7 ± 7.3	183.9 ± 7.8	183.0 ± 8.2	181.9 ± 5.2
RPE	7.2 ± 0.6	7.6 ± 0.7	7.6 ± 0.6	6.9 ± 0.3	6.9 ± 0.6

greater TD and DC at 14-17.9 km·h⁻¹ than without floaters, with the zone and square floaters (large and moderate magnitude) and greater DC at 7-13.9 km·h⁻¹ than without floaters or with the zone floaters (large and moderate magnitude). Comparing the changes in velocity, it was found that RPs achieved statistically more decelerations with square floaters than in any other format (large and moderate magnitude). With external floaters and without floaters, RPs performed more accelerations than with the internal zone floaters (large to small magnitude). In the format with internal floaters, RPs performed statistically fewer decelerations than with the

external floaters (moderate magnitude). The RPE of RPs when playing with internal and external floaters were statistically higher than in any other format of SSGs (moderate magnitude). RPs did not demonstrate statistical differences in the HR_{mean} and HR_{max} across any evaluated drill.

Comparisons of regular vs. floater players

Figure 4 reports the comparison of external and internal load between RPs and floaters. The analyses indicate that RPs achieved statistically greater



Bars indicate uncertainty in the true mean changes with 90% confidence intervals. Trivial (shaded) areas were calculated from the smallest worthwhile change. * TD = Total distance covered; DC = Distance covered; Acc = accelerations; Dec = decelerations; HR = heart rate; RPE = rate of perceived exertion; ES = Effect size; * $p < 0.05$; ** $p < 0.01$

Figure 4. Comparison of the external and internal load in RP and floater players in SSGs.

TD, DC at 7-13.9, 14-17.9, > 18 km h⁻¹, number of accelerations and RPE than floaters in all the SSG formats (very large to moderate magnitude). Regular players also reached a statistically greater number of decelerations and lower HR_{max} than the external, zone and square floaters, and a higher HR_{mean} than the zone and square floaters (very likely to moderate magnitude).

Discussion and conclusions

The main findings of this study were that the external and internal load of RPs and floaters were influenced by floaters' positions during the 4vs4+2, showing that (1) the floaters in the internal and external positions exhibited higher external and internal load than those in zone or square positions; (2) although internal and external floaters provoked greater RPE and external load in the RPs than did other floater positions or the absence of floaters, HR_{mean} and HR_{max} did not show difference in any format; and (3) independently of the floaters' positions, RPs had greater internal and external load than the floaters.

This study has shown that floaters have specific requirements depending on their position in the 4vs4+2. Our results revealed that the internal position was the most demanding scenario for floaters. Their freedom of movement throughout the available space could be the cause of this. The square and zone floaters showed lower internal (RPE and HR) and external (DC at 7-13.9, 14-17 and > 18 km·h⁻¹) load than the other positions. Their central position in the game may allow them to focus on adopting a suitable placement and body position to receive and pass the ball during the game, while internal and external floaters have to move constantly to receive the ball. Contrary to previous works (Casamichana & Castellano, 2010; Gaudino, Alberti, & Iaia, 2014; Owen, Wong, Paul, & Dellal, 2014), in this study a smaller relative area per player— zone (100m²) < internal (120 m²) < external and square (200 m²)— did not decrease the loads of the players, which could suggest that other aspects such as technical-tactical components of SSGs maybe have a more important influence on the behavior of the players. This information may be of value in the training of post-injury or over-trained players without leaving a specific context since floater training load can be manipulated by modifying their position. Taking this into account, and from a physical and physiological perspective, a suitable progression could be to successively incorporate such a player into 4vs4+2 as a floater in: square, zone, external and finally internal area.

This study could also suggest that the internal and external load of RPs are influenced by the position of floaters in 4vs4+2 SSG, creating inequalities in the magnitude of the training load and in its nature (TD, accelerations or decelerations). Each

floater position can modify the shape of the game and the position of RPs, affecting the demands imposed on them. Although in line with the findings of Casamichana, Bradley, and Castellano (2018), our results contrast with those obtained by Gollin et al. (2016), who found similarities in the running activity of RPs despite modifications in floater positions. These discrepancies could be due to the fact that Gollin et al. (2016) used a format with a relative area of approximately 80m² per player, while our study used 120 and 150 m² per player, as did Casamichana et al. (2018). This is especially important because it has been affirmed that player loads statistically increase in areas above 100 m² per player (Sarmiento, et al., 2018), so it could be suggested that floater position is less determinant in soccer drills with small relative areas than in those with larger areas. Further, our data are not in line with previous findings comparing RPs' activity in SSGs with distinct areas and with floaters (Lacome, et al., 2018). This is possibly because that study obtained greater RPs' demands in the format with a larger relative area, and only used SSGs with one floater. In addition, their players were professionals, and the difference in the relative area between their SSGs was approximately half. These circumstances may explain the conflicting results between the studies.

Another relevant aspect that could explain the disparities in the RPs' load in SSGs is that all players change their spatial dispersion, the way they play, and the distances they maintain from their direct opponents when floaters are manipulated in SSGs (Sampaio, et al., 2014). It is known that modifying the length of the pitch increases the load more than modifying its width (Casamichana, et al., 2018). This can also be achieved (without modifying the SSG space) by including internal and external floaters who increase the running activity and internal load of RPs, while with zone and square floaters RPs can prioritize the central region of the game where floaters are placed, allowing them to perform fewer movements to recover the ball. Responses from SSGs played without floaters are as expected, since it has been demonstrated that playing with small unbalanced situations can have a great impact on the demand of RPs (Torres-Ronda, et al., 2015). Despite the importance and the high incidence of floaters in SSGs during soccer training (Hill-Haas, et al., 2010; Lacome, et al., 2018; Rabano-Munoz, et al., 2019), this topic has not received much attention in literature (Ric, et al., 2015), so more research is needed to advance knowledge, and therefore, to be able to compare data with other studies.

Comparing the running activity and physiological demands of RPs and floaters during the SSGs, our results are in line with previous studies showing a greater external load in RPs (Gollin, et al., 2016; Lacome, et al., 2018; Rabano-Munoz, et al., 2019).

In relation to the internal load and depending on the floater position, our data are in line with Rabano-Munoz et al. (2019) that showed a greater internal load in RPs than in floaters, and with Lacombe et al. (2018) who did not report differences in HR between RPs and floaters. While our variations could be due to the dissimilar load of the floaters, the discrepancies between these studies may be due to different standard levels of players and the probable dissociation between HR and VO₂ during SSGs. We think that the fact that RPs have to participate in both phases of the game (offensive and defensive) created greater demands than those in the floaters, who only focused their activity on the offensive phase of the game. Using players as floaters during 4vs4+2 could be a simple way to decrease the external and internal load for optimal management in specific cases (e.g. during the return-to-play period after injury).

This study has some limitations that mean the data should be interpreted with caution. Only one format (4vs4+2) was used, with a standardized pitch size, so it is difficult to extrapolate these results to SSGs with dissimilar numbers of players or field sizes. In future studies the format and orientation of the floaters should be modified. In addition, this work was carried out with semi-professional players and the size of the sample was small, so it is unclear whether players of other standard levels or a greater sample would display the same behavior. It would therefore be interesting and useful to conduct

further comparative studies on the load of floaters and RPs in SSGs, varying the number of players, the pitch area and the level of the players.

This study showed that the incorporation of floaters during 4vs4+2 SSGs influenced the internal and external load of both the RPs and floaters. The floaters placed in the internal and external positions had higher running and physiological demands than those in the zone or square positions and provoked a greater internal and external load in RPs. Finally, RPs had always greater internal and external load than floaters, regardless of the 4vs4+2 format. Considering these conclusions, some practical applications may be offered:

1. Players could be used as floaters during 4vs4+2 to minimize their training load in some special circumstances (e.g., over-trained players or in a return-to-play process), allowing them to train with a lower training load than RPs.
2. Coaches could modify floater positions depending on the magnitude of the training load they want to apply, choosing the internal and external positions to increase it and the zone and square positions to decrease it.
3. Coaches could use floaters in their SSGs to provoke a greater internal and external load during training sessions, depending on players' roles in the team. For example, on the first day post-match starter players could be used as zone floaters to facilitate their active recovery.

References

- Aguiar, M., Botelho, G., Lago, C., Macas, V., & Sampaio, J. (2012). A review on the effects of soccer small-sided games. *Journal of Human Kinetics*, 33, 103-113. doi: 10.2478/v10078-012-0049-x
- Barbero-Alvarez, J.C., Coutts, A., Granda, J., Barbero-Alvarez, V., & Castagna, C. (2010). The validity and reliability of a global positioning satellite system device to assess speed and repeated sprint ability (RSA) in athletes. *Journal of Science and Medicine in Sport*, 13(2), 232-235. doi: 10.1016/j.jsams.2009.02.005
- Borg, G., Hassmen, P., & Lagerstrom, M. (1987). Perceived exertion related to heart rate and blood lactate during arm and leg exercise. *European Journal of Applied Physiology and Occupational Physiology*, 56(6), 679-685. doi: 10.1007/bf00424810
- Buchheit, M., Al Haddad, H., Simpson, B.M., Palazzi, D., Bourdon, P.C., Di Salvo, V., & Mendez-Villanueva, A. (2014). Monitoring accelerations with GPS in football: Time to slow down? *International Journal of Sports Physiology and Performance*, 9(3), 442-445. doi: 10.1123/ijsp.2013-0187
- Campos-Vazquez, M.A., Casamichana, D., Suarez-Arrones, L., González-Jurado, J.A., Toscano-Bendala, F.J., & León-Prados, J.A. (2017). Medium-sided games in soccer: Physical and heart rate demands throughout successive working periods. *Journal of Human Sports and Exercise*, 12(1), 129-141. doi: 10.14198/jhse.2017.121.11
- Casamichana, D., Bradley, P.S., & Castellano, J. (2018). Influence of the varied pitch shape on soccer players physiological responses and time-motion characteristics during small-sided games. *Journal of Human Kinetics*, 64, 171-180. doi: 10.1515/hukin-2017-0192
- Casamichana, D., & Castellano, J. (2010). Time-motion, heart rate, perceptual and motor behaviour demands in small-sides soccer games: Effects of pitch size. *Journal of Sports Sciences*, 28(14), 1615-1623. doi: 10.1080/02640414.2010.521168
- Casamichana, D., Castellano, J., & Castagna, C. (2012). Comparing the physical demands of friendly matches and small-sided games in semiprofessional soccer players. *Journal of Strength and Conditioning Research*, 26(3), 837-843. doi: 10.1519/JSC.0b013e31822a61cf

- Castillo, D., Raya-Gonzalez, J., Manuel Clemente, F., & Yanci, J. (2020). The influence of offside rule and pitch sizes on the youth soccer players' small-sided games external loads. *Research in Sports Medicine*, 28(3), 324-338. doi: 10.1080/15438627.2020.1739687
- Clemente, F.M., Afonso, J., Castillo, D., Arcos, A.L., Silva, A.F., & Sarmento, H. (2020). The effects of small-sided soccer games on tactical behavior and collective dynamics: A systematic review. *Chaos Solitons and Fractals*, 134, 109710.
- Clemente, F.M., Praca, G.M., Brecht, S., van der Linden, C.M.I., & Serra-Olivares, J. (2019). External load variations between medium- and large-sided soccer games: Ball possession games vs regular games with small goals. *Journal of Human Kinetics*, 70, 191-198. doi: 10.2478/hukin-2019-0031
- Coutinho, D., Goncalves, B., Travassos, B., Folgado, H., Figueira, B., & Sampaio, J. (2020). Different marks in the pitch constraint youth players' performances during football small-sided games. *Research Quarterly for Exercise and Sport*, 91(1), 15-23. doi: 10.1080/02701367.2019.1645938
- Coutts, A.J., & Duffield, R. (2010). Validity and reliability of GPS devices for measuring movement demands of team sports. *Journal of Science and Medicine in Sport*, 13(1), 133-135. doi: 10.1016/j.jsams.2008.09.015
- Coutts, A.J., Rampinini, E., Marcora, S.M., Castagna, C., & Impellizzeri, F.M. (2009). Heart rate and blood lactate correlates of perceived exertion during small-sided soccer games. *Journal of Science and Medicine in Sport*, 12(1), 79-84. doi: 10.1016/j.jsams.2007.08.005
- Davids, K., Araujo, D., Correia, V., & Vilar, L. (2013). How small-sided and conditioned games enhance acquisition of movement and decision-making skills. *Exercise and Sport Sciences Reviews*, 41(3), 154-161. doi: 10.1097/JES.0b013e318292f3ec
- Dellal, A., Chamari, K., Pintus, A., Girard, O., Cotte, T., & Keller, D. (2008). Heart rate responses during small-sided games and short intermittent running training in elite soccer players: A comparative study. *Journal of Strength and Conditioning Research*, 22(5), 1449-1457. doi: 10.1519/JSC.0b013e31817398c6
- Dellal, A., Owen, A., Wong, D.P., Krusturup, P., van Exsel, M., & Mallo, J. (2012). Technical and physical demands of small vs. large sided games in relation to playing position in elite soccer. *Human Movement Science*, 31(4), 957-969. doi: 10.1016/j.humov.2011.08.013
- Faulkner, J.A. (1968). *Physiology of swimming and diving*. Baltimore: Academic Press.
- Foster, C., Florhaug, J.A., Franklin, J., Gottschall, L., Hrovatin, L.A., Parker, S., . . ., & Dodge, C. (2001). A new approach to monitoring exercise training. *Journal of Strength and Conditioning Research*, 15(1), 109-115.
- Gaudino, P., Alberti, G., & Iaia, F.M. (2014). Estimated metabolic and mechanical demands during different small-sided games in elite soccer players. *Human Movement Science*, 36, 123-133. doi: 10.1016/j.humov.2014.05.006
- Gollin, M., Alfero, S., & Abate Daga, F. (2016). Manipulation of playing field's length/width ratio and neutral players' positioning: Activity profile and motor behavior demands during positional possession soccer small sided games in young elite soccer players. *International Journal of Sports Science*, 6(3), 106-115. doi: 10.5923/j.sports.20160603.07
- Hill-Haas, S.V., Coutts, A.J., Dawson, B.T., & Rowsell, G.J. (2010). Time-motion characteristics and physiological responses of small-sided games in elite youth players: The influence of player number and rule changes. *Journal of Strength and Conditioning Research*, 24(8), 2149-2156. doi: 10.1519/JSC.0b013e3181af5265
- Hill-Haas, S.V., Dawson, B., Impellizzeri, F.M., & Coutts, A.J. (2011). Physiology of small-sided games training in football: A systematic review. *Sports Medicine*, 41(3), 199-220. doi: 10.2165/11539740-000000000-00000
- Hill-Haas, S.V., Rowsell, G.J., Dawson, B.T., & Coutts, A.J. (2009). Acute physiological responses and time-motion characteristics of two small-sided training regimes in youth soccer players. *Journal of Strength and Conditioning Research*, 23(1), 111-115. doi: 10.1519/jsc.0b013e3181818efc1a
- Hopkins, W.G., Marshall, S.W., Batterham, A.M., & Hanin, J. (2009). Progressive statistics for studies in sports medicine and exercise science. *Medicine and Science in Sports and Exercise*, 41(1), 3-13. doi: 10.1249/MSS.0b013e3181818cb278
- Katis, A., & Kellis, E. (2009). Effects of small-sided games on physical conditioning and performance in young soccer players. *Journal of Sports Science and Medicine*, 8(3), 374-380.
- Kelly, D.M., & Drust, B. (2009). The effect of pitch dimensions on heart rate responses and technical demands of small-sided soccer games in elite players. *Journal of Science and Medicine in Sport*, 12(4), 475-479. doi: 10.1016/j.jsams.2008.01.010
- Lacome, M., Simpson, B.M., Cholley, Y., & Buchheit, M. (2018). Locomotor and heart rate responses of floaters during small-sided games in elite soccer players: Effect of pitch size and inclusion of goalkeepers. *International Journal of Sports Physiology and Performance*, 13(5), 668-671. doi: 10.1123/ijsp.2017-0340
- Low, B., Coutinho, D., Goncalves, B., Rein, R., Memmert, D., & Sampaio, J. (2020). A systematic review of collective tactical behaviours in football using positional data. *Sports Medicine*, 50(2), 343-385. doi: 10.1007/s40279-019-01194-7
- Mackenzie, R., & Cushion, C. (2013). Performance analysis in football: A critical review and implications for future research. *Journal of Sports Sciences*, 31(6), 639-676. doi: 10.1080/02640414.2012.746720
- Mallo, J., & Navarro, E. (2008). Physical load imposed on soccer players during small-sided training games. *Journal of Sports Medicine and Physical Fitness*, 48(2), 166-171.

- Owen, A.L., Newton, M., Shovlin, A., & Malone, S. (2020). The use of small-sided games as an aerobic fitness assessment supplement within elite level professional soccer. *Journal of Human Kinetics*, 71, 243-253. doi: 10.2478/hukin-2019-0086
- Owen, A.L., Wong, D.P., Paul, D., & Dellal, A. (2014). Physical and technical comparisons between various-sided games within professional soccer. *International Journal of Sports Medicine*, 35(4), 286-292. doi: 10.1055/s-0033-1351333
- Praca, G.M., Andrade, A.G.P., Brecht, S., Moura, F.A., & Moreira, P.E.D. (2022). Progression to the target vs. regular rules in soccer small-sided games. *Science and Medicine in Football*, 6(1), 66-71. doi: 10.1080/24733938.2020.1869811
- Rabano-Munoz, A., Asian-Clemente, J., Saez de Villarreal, E., Naylor, J., & Requena, B. (2019). Age-related differences in the physical and physiological demands during small-sided games with floaters. *Sports (Basel)*, 7(4). doi: 10.3390/sports7040079
- Rampinini, E., Impellizzeri, F.M., Castagna, C., Abt, G., Chamari, K., Sassi, A., & Marcora, S.M. (2007). Factors influencing physiological responses to small-sided soccer games. *Journal of Sports Sciences*, 25(6), 659-666. doi: 10.1080/02640410600811858
- Ric, A., Hristovski, R., & Torrents, C. (2015). Can joker players favor the exploratory behaviour in football small-sided games? *Research in Physical Education, Sport and Health*, 4(2), 35-39.
- Rico-Gonzalez, M., Pino-Ortega, J., Praca, G. M., & Clemente, F. M. M. (2022). Practical applications for designing soccer training tasks from multivariate data analysis: A Systematic review emphasizing tactical training. *Perceptual and Motor Skills*, 315125211073404. doi: 10.1177/00315125211073404
- Sampaio, J., Lago, C., Goncalves, B., Macas, V.M., & Leite, N. (2014). Effects of pacing, status and unbalance in time motion variables, heart rate and tactical behaviour when playing 5-a-side football small-sided games. *Journal of Science and Medicine in Sport*, 17(2), 229-233. doi: 10.1016/j.jsams.2013.04.005
- Sanchez-Sanchez, J., Hernandez, D., Casamichana, D., Martinez-Salazar, C., Ramirez-Campillo, R., & Sampaio, J. (2017). Heart rate, technical performance, and session-RPE in elite youth soccer small-sided games played with wildcard players. *Journal of Strength and Conditioning Research*, 31(10), 2678-2685. doi: 10.1519/JSC.0000000000001736
- Sarmento, H., Clemente, F.M., Harper, L.D., Costa, I.T., Owen, A., & Figueiredo, A.J. (2018). Small sided games in soccer—A systematic review. *International Journal of Performance Analysis in Sport*, 18(5), 693-749. doi: https://doi.org/10.1080/24748668.2018.1517288
- Suarez-Arrones, L., Nunez, J., Saez de Villarreal, E., Galvez, J., Suarez-Sanchez, G., & Munguia-Izquierdo, D. (2016). Repeated-high-intensity-running activity and internal training load of elite rugby sevens players during international matches: A comparison between halves. *International Journal of Sports Physiology and Performance*, 11(4), 495-499. doi: 10.1123/ijsp.2014-0523
- Torres-Ronda, L., Goncalves, B., Marcelino, R., Torrents, C., Vicente, E., & Sampaio, J. (2015). Heart rate, time-motion, and body impacts when changing the number of teammates and opponents in soccer small-sided games. *Journal of Strength and Conditioning Research*, 29(10), 2723-2730. doi: 10.1519/JSC.0000000000000932
- Turner, A.N., & Stewart, P.F. (2014). Strength and conditioning for soccer players. *Strength and Conditioning Journal*, 36(4), 1-13.
- Vilar, L., Araujo, D., Davids, K., & Button, C. (2012). The role of ecological dynamics in analysing performance in team sports. *Sports Medicine*, 42(1), 1-10. doi: 10.2165/11596520-000000000-00000
- Yue, Z., Broich, H., Seifriz, F., & Mester, J. (2008). Mathematical analysis of a soccer game. Part I: Individual and collective behaviors. *Studies in Applied Mathematics*, 121(3), 223-243. doi: 10.1111/j.1467-9590.2008.00413.x

Submitted: August 24, 2021

Accepted: March 4, 2022

Published Online First: May 9, 2022

Correspondence to:

Jose Antonio Asian-Clemente, Ph.D.

University Pablo de Olavide,

Department of Sport Sciences,

Ctra. Utrera 1, 41500 Seville, Spain

Phone: +34 954977601

E – mail: jasicle@upo.es