

TRAINING LOAD AND PLAYERS' READINESS MONITORING METHODS USED IN VOLLEYBALL: A SYSTEMATIC REVIEW

Roberto Vavassori¹, M. Perla Moreno Arroyo¹, and Aurelio Ureña Espa^{1,2}

¹Department of Sports Science and Physical Education, University of Granada, Granada, Spain

²University Mixed Institute of Sport and Health iMUDS, University of Granada, Granada, Spain

Review

DOI 10.26582/k.56.1.10

Abstract:

Monitoring workloads during training and competition and players' readiness seems to be key to increasing performance, reducing injury incidence and avoiding overtraining. We systematically reviewed the methods used to measure workloads and athletes' readiness in volleyball to help coaches make the best decision when selecting monitoring methods. Databases Web of Science, Scopus, SPORTDiscus and PubMed were searched from inception to the 21st of February 2022. All peer-reviewed original research in English, Spanish, Portuguese and Italian, longitudinally monitoring loads and athlete readiness in indoor volleyball team settings of any level, gender and age were included. The quality of evidence was evaluated with a modified risk of bias assessment used in previous research by Castellano et al. (2014). This study has been registered in PROSPERO ID CRD42022316313. Out of 1774 records identified, 78 were screened of which 55 full texts were added for systematic review. For internal workload, the session rating of perceived exertion (sRPE) seems to be the "golden standard" used from 2010 to 2022 across all the studies. External workload has mainly been researched through quantified jumps. Even with technological advances and the introduction of microsensors in 2017, the use of video analysis is still present nowadays. Players' readiness studies mainly used the total quality recovery scale (TQR) and wellness questionnaires in most research. New technological advances offer coaches more extensive and real-time data on external load. However, the use of the sRPE, TQR/WB, CMJ would create a monitoring system sufficient for teams at developmental stages and are of a reduced cost.

Keywords: workload, monitoring methods, wellness, readiness, performance

Introduction

Volleyball is a dynamic and unpredictable sport that stands out for the combination of high-intensity efforts with short periods of rest at low intensity. Among the skills that a volleyball player should possess, the following stand out: at a physical level, lower limb power, accelerations and decelerations over short distances (Sheppard & Newton, 2012), and at a technical level: setting, serving, blocking and attacking, which are highly influenced by the jumping action (Sheppard, Nolan & Newton, 2012).

Due to the high density of eccentric actions, together with the high number of impacts generated by landings and braking, an increase in fatigue and muscle damage is to be expected, which can lead to a decrease in athletic performance (Eliakim, et al., 2009; Souglis, Bogdanis, Giannopoulou, Pappadopoulos & Apostolidis, 2015). That, in addition to the competitive density of the sport itself and

certain contextual factors, can lead to a suboptimal recovery state (Clemente, et al., 2017; Fessi, et al., 2016).

This highlights the importance of knowing the state of our athletes and their progression towards previously established objectives. Also, that knowledge helps in the decision-making of coaches and technical staff regarding possible modifications in planning (Jeffries, et al., 2022). To this end, it seems essential to know the effects of training and competition on athletes at physiological, psychological and biomechanical levels, among others. More specifically, it is necessary to analyze training effects from the point of view of the work performed (external load) and of the response of athletes' body to the performance of this work (internal load). Coaches need to know the effect of the loads applied (acute, chronic, positive and negative) and the contextual and individual factors of each player (Jeffries, et al., 2022).

The range of load quantification methods that have been used over the years is very extensive but we can observe that certain methods stand out in scientific literature over others. Among the internal load quantification methods, the subjective perception of effort in the session (sRPE) (Foster, et al., 2001) or, at the objective level, heart rate and training impulse (TRIMP). Among the most current external load quantification methods are GPS systems, microsensors, and accelerometers (Bourdon, et al., 2017). It is also worth mentioning those tools that allow us to know the state of athletes' readiness through questionnaires (wellness or wellbeing). Although not scientifically validated (Jeffries, et al., 2020), they are widely used by sports professionals in their decision-making and can be at the same level of use as the total quality recovery scale (TQR) or the recovery-stress questionnaire for athletes (REST-Q) scales.

Despite a large number of options available, a consensus on the methods that should be used for monitoring athletes (Scott, Duthie, Thornton & Dascombe, 2016) does not yet exist and, specifically in volleyball, there is a lack of clarity on the tools that should be used to monitor loads and players' readiness status. Although some recent reviews have been able to detect the tools used in volleyball (Pisa, Zecchin, Gomes, Norberto & Puggina, 2022), they have only focused on internal loads with professional and male players, which could still generate a total lack of clarity. Therefore, this study aims to systematically review the scientific literature to know the methods and tools used in volleyball for the control of training loads, match loads and readiness status in volleyball teams with the secondary objective of helping coaches and technical staff in the decision-making process when selecting the most appropriate monitoring and readiness tools for their teams.

Methods

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 protocol was used for this systematic review (Page, et al., 2021).

Research strategy

A systematic search of four electronic databases was conducted: PubMed, Scopus, Web of Science, and SPORTDiscus. The combination of different terms in title, abstract or keywords was made as follows: (Volleyball AND ("monitor*" OR "control" OR "record*" OR "quantif*") AND ("load*" OR "internal load*" OR "external load*" OR "training load*" OR "match load*" OR "internal training load*" OR "external training load*" OR "workload" OR "training intens*" OR "training respon*" OR "subjective" OR "objective" OR "fatigue" OR

"non-functional overreaching" OR "recovery" OR "readiness" OR "wellness" OR "wellness questionnaire" OR "wellbeing" OR "well being" OR "well-being" OR "mood" OR "stress" OR "sleep") NOT "beach volleyball"). The range for years was established from the earliest available record to the 21st of February 2022. To reduce the chances of studies being left out, reference lists of included articles and relevant reviews were scanned to ensure a wider reach of our search.

Inclusion and exclusion criteria

Eligibility criteria were established using the PICO model from the PRISMA 2020 report (Page, et al., 2021):

The manuscripts selected in this systematic review followed these criteria: (1) studies based on either internal load, external load, readiness for training/matches or any combination of the three; (2) studies collecting longitudinal data of workloads and/or player's readiness in training, matches or both events; (3) articles on indoor volleyball; (4) English, Italian, Portuguese or Spanish versions of the studies; (5) original research published in a peer-reviewed journal of players enrolled in a team setting of any age, level or gender; (6) studies from the database inception to 21st of February 2022.

Exclusion criteria were established as follows: (1) studies evaluating injury prevention or reduction; (2) studies on beach volleyball; (3) manuscripts checking validity and reliability or focusing on specific drills or testing specific individual physiological demands; (4) studies with exact measurements (5) experimental studies, conference abstracts or unpublished manuscripts.

Study selection

Database search results were added to reference manager Mendeley (Elsevier, London, UK) where duplicate articles were removed. Titles and abstracts screening of remaining records was performed by the first author RV. Then, full texts were analyzed against inclusion criteria by RV and, in case of uncertainty, MPMA and AUE were consulted for discussion and reaching a final consensus.

Data collection

The first author examined and extracted information from the selected studies to be included in the systematic review into a specifically created spreadsheet. When possible, the following data were extracted from each article following the "Population, Intervention, Comparison, Outcome" (PICO) framework: (1) sample size, gender, age, playing level and country; (2) study duration, study period, study observation (only training, only matches or both); (3) instruments used (e.g. sRPE, questionnaires, video analysis), characteristics of instru-

ments (scales, devices, thresholds); (4) study goals, study variables, main results, outcome and conclusions; (5) statistical analysis; (6) study design.

Risk-of-bias assessment

Studies were evaluated qualitatively using modified assessment criteria from Castellano, Alvarez-Pastor, and Bradley (2014) (Table 1). The main modifications were as follows: item eight was removed from the original tool as irrelevant to the current review, and answers in item 7 were converted into "YES" or "NO", so all the questions could be affirmative or negative to avoid question scores. Finally, rewording of the remaining eight items was applied to better adapt the tool to this systematic review criteria. A maximum of eight positive responses could be achieved depending on how the criteria were met. The risk-of-bias assessment was used to weigh a study's contribution to the results. Articles with a positive response of five or above were considered to carry full weight, whereas for those with four or fewer "Yes" contributions to the results were halved. RV applied the tool to each of the included studies; in the case of discrepancies, they were solved by a discussion with the remaining authors.

Data synthesis

The synthesis of data was made descriptively with the information presented in text and detailed tables. The goal of this systematic review was to observe the most used methods for monitoring workloads and players' readiness in volleyball. Since a recompilation of the results of the studies was not sought, meta-analysis was not taken into consideration. The main goal of a meta-analysis is to statistically analyze results from a relatively homogeneous group of studies, to integrate their results. The selected studies were deemed heterogeneous in variables, methods, interventions, reporting, outcome measures and study designs. Also, meta-analysis can only analyze studies with specific statistical information, therefore discarding qualitative studies.

Results

Selected studies

Initially, 1774 records were retrieved from the different databases (PubMed = 307, SPORTDiscus = 423, Web of Science = 443, Scopus = 601). A total of 677 were removed as duplicates. After screening the remaining titles and abstracts, 78 articles were selected for the full-text analysis. The rationale for rejecting full texts was as follows: language (Çelebi & Aksu, 2018; Maksimenko, Maksimenko, Zhilina & Bayeva, 2019; Sattler, 2021), not considered monitoring or athletes' readiness research (Bara Filho, de Andrade, Nogueira & Nakamura, 2013; Garcia-de-Alcaraz, Valadés & Palao, 2017; Mroczek, et al., 2014; Podstawski, Boraczynski, Nowosielska-Swadzba & Zwolinska, 2014; Zhou, 2021), not on volleyball (Hamlin, Wilkes, Elliot, Lizamore & Kathiravel, 2019), injuries and/or rehabilitation studies (Hurd, Hunter-Giordano, Axe & Snyder-Mackler, 2009; Sole, Kavanaugh & Stone, 2017; Visnes & Bahr, 2013), not following up continuously or longitudinally (Hank, Zahalka & Maly, 2015; Horta, Bara Filho, Miranda, Coimbra & Werneck, 2017; Hurd, et al., 2009; Moreira, et al., 2013; Mortatti, Pinto, Lambertucci, Hirabara & Moreira, 2018; Noce, et al., 2008; Pires & Ugrinowitsch, 2016; Pires & Ugrinowitsch, 2021; Reynoso-Sánchez, et al., 2016) and no full-text available due to journal's embargo (Gielen, Mehuys, Berckmans, Meeusen & Aerts, 2022; Ungureanu, Brustio, Boccia, Rainoldi & Lupo, 2021; Xue, 2017). After further exploration of references, two records were recouped (Freitas, Miloski, & Bara Filho, 2015; Lacerda, et al., 2015) and added to the final 55 studies included in the systematic review. (See Table 2 for reference of the included studies.) A flow chart of the process is presented in Figure 1.

Studies were published between 2010 and 2022. Out of the 55 studies included in this review, 10 studies covered the 2010-2015 period, 33 period of 2016-2020 and 12 two years 2021-2022. Regarding gender, 33 studies included only male participants, 19 female, and three both. In terms of level, 32

Table 1. Risk of bias assessment criteria

Criteria	Answer
1 The study is published in a peer-reviewed journal	No Yes
2 The study is published in an indexed journal	No Yes
3 The study objective(s) is/are clearly set out	No Yes
4 Either the number of recordings is specified or the distribution of player/recordings used is known	No Yes
5 The duration of player recordings (how many weeks/training sessions, how many matches, etc.)	No Yes
6 A distinction is made according to player position, training session type and/or match	No Yes
7 The reliability/validity of the instrument is not stated or is mentioned	No Yes
8 The results are clearly presented	No Yes




Fig. 1. Studies selection process flow chart recommended in the PRISMA, 2020. Outlining the path followed to select articles included in the systematic review

Table 2. Participants' characteristics (number, gender, level) and results of methodological quality assessment of a study

Study	Year	N	Gender	Level	Quality questions								Total
					Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	
Andrade et al.	2021	15	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
Aoki et al.	2017	18	Male	U16 & U19	Y	Y	Y	N	N	Y	Y	Y	6
Bahr & Bahr	2014	44	Both	Junior volleyball Norway	Y	Y	Y	Y	Y	Y	N	Y	7
Brandão et al.	2018	14	Male	Professional	Y	N	Y	Y	Y	Y	Y	Y	7
Cardoso et al.	2021	9	male	Professional Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
Carroll et al.	2019	11	Female	NCAA D1	Y	Y	Y	Y	Y	Y	N	Y	7
Castello et al.	2018	10	Female	NCAA D1	Y	N	Y	Y	N	N	Y	Y	5
Clemente et al.	2019	13	Male	Professional Portugal	Y	Y	Y	Y	Y	Y	Y	Y	8
Clemente et al.	2020	13	Male	Professional Portugal	Y	Y	Y	Y	Y	Y	N	Y	7
Coyne et al.	2021	63	Female	Olympic level	Y	Y	Y	Y	Y	Y	Y	Y	8
de Andrade et al.	2014	15	Male	National level Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
De Leeuw et al.	2021	10	Male	Elite	Y	Y	Y	Y	Y	Y	Y	Y	8
Debien et al.	2018	15	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
Duarte et al.	2019	14	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
Duarte et al.	2019	15	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
Edmonds, Schmidt & Siedlik	2021	14	Female	NCAA D1	Y	Y	Y	Y	Y	Y	Y	Y	8
Freitas et al.	2014	16	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	Y	Y	8
Freitas et al.	2015	7	Male	Under 16 Brazil	Y	Y	Y	Y	Y	Y	Y	Y	8
Freitas, Miloski & Bara Filho	2015	12	Male	National league	Y	Y	Y	Y	Y	Y	N	Y	7
García-de-Alcaraz et al.	2020	11	Male	Professional Spain	Y	Y	Y	Y	Y	Y	N	Y	7
Háp et al.	2011	8	Male	Professional Czech Republic	Y	Y	Y	Y	Y	Y	N	Y	7
Haraldsdottir et al.	2021	17	Female	NCAA D1	Y	Y	Y	N	Y	N	N	Y	5

Table 2. (Continued)

Study	Year	N	Gender	Level	Quality questions								Total
					Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	
Hernández-Cruz et al.	2017	12	Male	Professional Mexico	Y	Y	Y	N	Y	Y	Y	Y	7
Herring & Fukuda	2022	14	Female	NCAA Div 1	Y	Y	Y	Y	Y	Y	Y	Y	8
Horta et al.	2017	15	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	Y	Y	8
Horta et al.	2019	12	male	elite	Y	Y	Y	Y	Y	Y	Y	Y	8
Horta et al.	2019	12	Male	Professional Brazil	Y	Y	Y	Y	N	Y	N	Y	6
Horta et al.	2020	9	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	Y	Y	8
Kraft et al.	2020	56	Female	NCAA D2	Y	Y	Y	Y	Y	Y	Y	Y	8
Kupperman et al.	2021	11	Female	NCAA Div 1	Y	Y	Y	Y	N	Y	N	Y	6
Lacerda et al.	2015	8	male	professional	Y	Y	Y	Y	Y	Y	N	Y	7
Libs et al.	2019	3	Female	NCAA D1	Y	Y	Y	Y	Y	Y	N	Y	7
Lima et al.	2019	5	Male	Professional Portugal	Y	Y	Y	Y	Y	Y	Y	Y	8
Lima et al.	2020	8	Male	Professional Portugal	Y	Y	Y	Y	Y	Y	Y	Y	8
Lima et al.	2021	10	Male	Portuguese 1st division	Y	Y	Y	Y	Y	Y	Y	Y	8
Malisoux et al.	2013	269	both	Elite juniors	Y	Y	Y	Y	Y	Y	N	Y	7
Mendes et al.	2018	13	Male	Professional Portugal	Y	Y	Y	N	Y	Y	N	Y	6
Moreira et al.	2010	20	male	Juniors	Y	Y	Y	Y	Y	Y	N	Y	7
Piatti et al.	2021	12	Male	Elite	Y	Y	Y	Y	Y	Y	Y	Y	8
Rabbani et al.	2021	13	Female	Iran national team	Y	Y	Y	Y	Y	Y	N	Y	7
Rabello et al.	2019	18	Male	Top Dutch division	Y	Y	Y	Y	Y	Y	Y	Y	8
Rodríguez-Marroyo et al.	2014	12	Female	Spanish Primera National	Y	Y	Y	N	Y	Y	N	Y	6
Roy et al.	2019	15	Female	University Canada	Y	N	Y	N	Y	N	Y	Y	5
Roy et al.	2020	15	Female	University Canada	Y	N	Y	N	Y	N	Y	Y	5
Sanders et al.	2018	1	Female	NCAA D1	Y	Y	Y	Y	Y	Y	Y	Y	8
Skazalski et al.	2018	14	Male	Professional Qatar	Y	Y	Y	Y	Y	Y	Y	Y	8
Tavares et al.	2018	13	Male	U19 Portugal	Y	Y	Y	Y	Y	Y	Y	Y	8
Taylor et al.	2019	14	Female	NCAA D1	Y	Y	Y	Y	Y	Y	N	Y	7
Taylor et al.	2022	16	female	NCAA D1	Y	Y	Y	Y	Y	Y	Y	Y	8
Timoteo et al.	2017	12	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
Timoteo et al.	2021	14	Male	Professional Brazil	Y	Y	Y	Y	Y	Y	N	Y	7
Ungureanu et al.	2021	10	Female	Professional Italy	Y	Y	Y	Y	Y	Y	Y	Y	8
van der Does et al.	2017	86	Both	University Netherlands	Y	Y	Y	Y	Y	Y	Y	Y	8
Vlantes & Readdy	2017	11	Female	NCAA D1	Y	Y	Y	Y	Y	Y	Y	Y	8
Wolfe et al.	2019	19	Female	NCAA D1	Y	Y	Y	Y	N	Y	N	Y	6
Average												7,2	

NCAA: National Collegiate Athletic Association

Q1-Q8: Y= yes; N= no

articles focused on the professional and elite level, 14 on the university competition level and nine on juniors and recreational players.

Quality of the studies

The quality of the included studies was considered medium-high as an average of seven positive responses ("YES") were obtained and no study received less than five. This means all studies got the same weight for the results. A more explanatory description of quality is illustrated in Table 2.

Monitoring methods

Studies showed a tendency to use a combination of different methods (36 articles). However, it is important to point out that, from the remaining 19 studies using a single metric, the majority of them were able to retrieve more than one derivative from one method (e.g., microensors obtaining jump count, jump height, jumps per position, jump frequency), hence multiple metrics were obtained. From these articles, seven studies only used internal measures (Castello, Reed, Lund, & Mack, 2018;

de Andrade, et al., 2014; Freitas, Miloski, et al., 2015; Háp, et al., 2011; Horta, Coimbra, Miranda, Werneck, & Bara Filho, 2017; Horta, Bara Filho, Coimbra, Werneck, & Miranda, 2019; Malisoux, Frisch, Urhausen, Seil, & Theisen, 2013), eight exclusively external (Bahr & Bahr, 2014; Herring & Fukuda, 2022; Lima, Palao, Castro, & Clemente, 2019; Piatti, et al., 2021; Skazalski, Whiteley, & Bahr, 2018; Taylor, Kantor, Hockenjos, Barnes, & Dischiavi, 2019; Taylor, Barnes, Gombatto, Greenwood, & Ford, 2022; Wolfe, et al., 2019) and the remaining four investigated readiness (Carroll, Wagle, Sole, & Stone, 2019; Haraldsdottir, Sanfilippo, McKay, & Watson, 2021; Hernández-Cruz, et al., 2017; van der Does, Sanne Brink, Ardi Otter, Visscher, & Plechelmus Marie Lemmink, 2017). From studies combining measures, 22 mixed internal workload and readiness (Andrade, Fernandes, Miranda, Reis Coimbra, & Bara Filho, 2021; Cardoso, Berriel, Schons, Costa, & Kruel, 2021; Carroll, et al., 2019; Clemente, et al., 2019; de Andrade, et al., 2014; Duarte, Alves, et al., 2019; Edmonds, Schmidt, & Siedlik, 2021; Freitas, Nakamura, Miloski, Samulski, & Bara Filho, 2014; Freitas, Nakamura, et al., 2015; Herring & Fukuda, 2022; Lima, et al., 2021; Lima, et al., 2019; Malisoux, et al., 2013; Rabbani, Agha-Alinejad, Gharakhanlou, Rabbani, & Flatt, 2021; Rabello, Zwerver, Stewart, van den Akker-Scheek, & Brink, 2019; Skazalski, et al., 2018; Tavares, Simões, Matos, Smith, & Driller, 2018; Taylor et al., 2022; Timoteo, et al., 2021; Ungureanu, Lupo, Boccia, & Brustio, 2021), four internal and external loads with readiness (all 3 together) (Cardoso, et al., 2021; de Leeuw, van der Zwaard, van Baar, & Knobbe, 2022; Kupperman, Curtis, Saliba, & Hertel, 2021; Ungureanu, Lupo, et al., 2021), five internal and external loads (Libs, Boos, Shipley, Peacock, & Sanders, 2019; Lima, et al., 2021; Lima, Silva, Afonso, Castro, & Clem-

ente, 2020; Rabello, et al., 2019; Vlantes & Readdy, 2017), three used two different internal load measures (Duarte, Coimbra, et al., 2019; Rodríguez-Marroyo, Medina, García-López, García-Tormo, & Foster, 2014; Roy, Caya, Charron, Comtois, & Sercia, 2020) and other two different external load measures (Garcia-de-Alcaraz, Ramírez-Campillo, Rivera-Rodríguez, & Romero Moraleda, 2020; Sanders, Boos, Shipley, Sheadler, & Peacock, 2018). Following a timeline, we can observe 12 studies combining methods from 2010 to 2018 and then an exponential increase between 2019 to 2021 with 23 studies in this period.

A full descriptive illustration of monitoring measures in chronological order, to observe the evolution of methods through time, is available in Table 3 and Figure 2.

Internal load

Internal load was tracked in 41 studies (74.5%) (Andrade, et al., 2021; Aoki, et al., 2017; Brandão, et al., 2019; Cardoso, et al., 2021; Castello, et al., 2018; Clemente, et al., 2019, 2020; Coyne, Coutts, Newton, & Haff, 2021; de Andrade, et al., 2014; de Leeuw, et al., 2022; Debien, et al., 2018; Duarte, Alves, et al., 2019; Duarte, Coimbra, et al., 2019; Edmonds, et al., 2021; Freitas, et al., 2014; Freitas, Miloski, et al., 2015; Freitas, Nakamura, et al., 2015; Háp, et al., 2011; Horta, Coimbra, et al., 2017; Horta, Bara Filho, Coimbra, Miranda, & Werneck, 2019; Horta, Bara Filho, Coimbra, Werneck, et al., 2019; Horta, et al., 2020; Kraft, et al., 2020; Kupperman, et al., 2021; Lacerda, et al., 2015; Libs, et al., 2019; Lima, et al., 2021; Lima, et al., 2020; Malisoux, et al., 2013; Mendes, et al., 2018; Moreira, de Freitas, Nakamura, & Aoki, 2010; Rabbani, et al., 2021; Rabello, et al., 2019; Rodríguez-Marroyo, et al., 2014; Roy, et al., 2019, 2020; Tavares, et al., 2018; Timoteo, et al., 2017, 2021; Ungureanu,

Table 3. Characteristics of study duration and methods used to monitor load in each article

Study	Monitoring method	Year
Moreira et al.	sRPE ^a / RPE ^b / sRPE derivatives + Other readiness	2010
Háp et al.	Other internal	2011
Malisoux et al.	sRPE / RPE / sRPE derivatives	2013
Bahr & Bahr	Video analysis (Jump/Swing count/load)	2014
Freitas et al.	sRPE / RPE / sRPE derivatives + other internal + CMJ ^c /SJ ^d /Rsi ^e + TQR ^f + REST-Q ^g	2014
Rodríguez-Marroyo et al.	sRPE / RPE / sRPE derivatives + HR ^h	2014
de Andrade et al.	sRPE / RPE / sRPE derivatives	2014
Lacerda et al.	sRPE / RPE / sRPE derivatives + TQR	2015
Freitas et al.	sRPE / RPE / sRPE derivatives + CMJ/SJ/Rsi + REST-Q	2015
Freitas, Miloski & Bara Filho	sRPE / RPE / sRPE derivatives	2015
Vlantes & Readdy	sRPE / RPE / sRPE derivatives + Microsensor (Jump/swing count/load) (Jump/swing count/load)	2017
Timoteo et al.	sRPE / RPE / sRPE derivatives + TQR + WB	2017

Table 3. Characteristics of study duration and method used to monitor load in each article (continuation)

Study	Monitoring method	Year
Horta et al.	sRPE / RPE / sRPE derivatives	2017
Aoki et al.	sRPE / RPE / sRPE derivatives + other readiness	2017
Hernández-Cruz et al.	HRV ⁱ	2017
van der Does et al.	REST-Q	2017
Sanders et al.	Microsensor (Jump/swing count/load)	2018
Skazalski et al.	Microsensor (Jump/swing count/load)	2018
Brandão et al.	sRPE / RPE / sRPE derivatives + WB + TQR	2018
Tavares et al.	sRPE / RPE / sRPE derivatives + CMJ/SJ/Rsi + WB + other readiness	2018
Mendes et al.	sRPE / RPE / sRPE derivatives + WB	2018
Debien et al.	sRPE / RPE / sRPE derivatives + TQR	2018
Castello et al.	sRPE / RPE / sRPE derivatives	2018
Rabello et al.	sRPE / RPE / sRPE derivatives + Microsensor (Jump/swing count/load) + Video analysis (Jump/Swing count/load)	2019
Libs et al.	HR + Microsensor (Jump/swing count/load)	2019
Lima et al.	Microsensor (Jump/swing count/load)	2019
Wolfe et al.	Video analysis (Jump/Swing count/load) + Other external	2019
Taylor et al.	Video analysis (Jump/Swing count/load)	2019
Duarte et al.	sRPE / RPE / sRPE derivatives + TQR + WB	2019
Clemente et al.	sRPE / RPE / sRPE derivatives + WB	2019
Roy et al.	sRPE / RPE / sRPE derivatives + WB	2019
Duarte et al.	sRPE / RPE / sRPE derivatives + HR	2019
Horta et al.	sRPE / RPE / sRPE derivatives + other internal + REST-Q	2019
Horta et al.	sRPE / RPE / sRPE derivatives	2019
Carroll et al.	CMJ/SJ/Rsi	2019
Lima et al.	sRPE / RPE / sRPE derivatives + Microsensor (Jump/swing count/load)	2020
García-de-Alcaraz et al.	Video analysis (Jump/Swing count/load)	2020
Clemente et al.	sRPE / RPE / sRPE derivatives + WB	2020
Horta et al.	sRPE / RPE / sRPE derivatives + TQR + other readiness	2020
Kraft et al.	sRPE / RPE / sRPE derivatives + HR + other readiness	2020
Roy et al.	sRPE / RPE / sRPE derivatives + Other Internal	2020
Lima et al.	sRPE / RPE / sRPE derivatives + Microsensor (Jump/swing count/load) + Video analysis (Jump/Swing count/load) + other external	2021
Kupperman et al.	sRPE / RPE / sRPE derivatives + Microsensor (Jump/swing count/load) + WB + other external	2021
Piatti et al.	Microsensor (Jump/swing count/load)	2021
Ungureanu et al.,	sRPE / RPE / sRPE derivatives + HR + WB + Video analysis (Jump/Swing count/load)	2021
Rabbani et al.	sRPE / RPE / sRPE derivatives + HR + HRV + CMJ/SJ/Rsi + WB	2021
De Leeuw et al.	sRPE / RPE / sRPE derivatives + other external + WB	2021
Edmonds, Schmidt & Siedlik	HR + HRV + WB	2021
Haraldsdottir et al.	WB	2021
Andrade et al.	sRPE / RPE / sRPE derivatives + TQR	2021
Timoteo et al.	sRPE / RPE / sRPE derivatives + TQR	2021
Cardoso et al.,	sRPE / RPE / sRPE derivatives + HRV + TQR + other readiness	2021
Coyne et al.	sRPE / RPE / sRPE derivatives + other readiness	2021
Herring & Fukuda	Microsensor (Jump/swing count/load)	2022
Taylor et al.	Microsensor (Jump/swing count/load)	2022

Note. ^asRPE: session rating of perceived effort; ^bRPE: rating of perceived effort; ^cCMJ: counter movement jump; ^dSJ: squat jump ^eRSI: reactive strength index; ^fTQR: total quality recovery scale; ^gREST-Q: recovery-stress questionnaire; ^hHR: heart rate; ⁱHRV: heart rate variability.




Fig. 2. Chronological evolution of the use of different measurement instruments found in the selected studies included in this systematic review.

Lupo, et al., 2021; Vlantes & Readdy, 2017), 38 of them (92.7%) operated with the sRPE/RPE as a measure for internal training and competition loads (Andrade, et al., 2021; Aoki, et al., 2017; Brandão, et al., 2019; Cardoso, et al., 2021; Castello, et al., 2018; Clemente, et al., 2019, 2020; Coyne, et al., 2021; de Andrade, et al., 2014; de Leeuw, et al., 2022; Debien, et al., 2018; Duarte, Coimbra, et al., 2019; Duarte, Alves, et al., 2019; Freitas, et al., 2014; Freitas, Miloski, et al., 2015; Freitas, Nakamura, et al., 2015; Horta, Coimbra, et al., 2017; Horta, Bara Filho, Coimbra, Miranda, et al., 2019; Horta, Bara Filho, Coimbra, Werneck, et al., 2019; Horta, et al., 2020; Kraft, et al., 2020; Kupperman, et al., 2021; Lacerda, et al., 2015; Lima, et al., 2021; Lima et al., 2020; Malisoux, et al., 2013; Mendes, et al., 2018; Moreira, et al., 2010; Rabbani, et al., 2021; Rabello, et al., 2019; Rodríguez-Marroyo, et al., 2014; Roy, et al., 2019, 2020; Tavares, et al., 2018; Timoteo, et al., 2017, 2021; Ungureanu, Lupo, et al., 2021; Vlantes & Readdy, 2017). All the studies recording sRPE applied the Category Ratio Scale 10 (CR-10) (Foster, et al., 2001). Multiple studies (18) took advantage of the versatility of the sRPE using derivatives. The sum of daily workloads into a weekly internal training load (WITL), monotony, strain and acute to chronic workload ratio (ACWR) were the most employed (Andrade, et al., 2021; Clemente, et al., 2019, 2020; de Leeuw, et al., 2022; Debien, et al., 2018; Duarte, Coimbra, et al., 2019; Freitas, et al., 2014; Freitas, Miloski, et al., 2015; Horta, Coimbra, et al., 2017; Horta, Bara Filho, Coimbra, Werneck, et al., 2019; Horta, Bara Filho, Coimbra, Miranda, et al., 2019; Horta, et al., 2020; Lacerda, et al., 2015; Malisoux, et al., 2013; Rabbani, et al., 2021; Rodríguez-Marroyo, et al., 2014; Timoteo, et al., 2021). Objective internal measures were less

used among the selected studies, with seven records using HR (Duarte, Alves, et al., 2019; Edmonds, et al., 2021; Kraft, et al., 2020; Libs, et al., 2019; Rabbani, et al., 2021; Rodríguez-Marroyo, et al., 2014; Ungureanu, Lupo, et al., 2021) and three other methods such as: saliva and blood markers (Háp, et al., 2011; Horta, Bara Filho, Coimbra, Miranda, et al., 2019; Roy, et al., 2020). See Table 3.

Regarding the usage of the above measures through the years, the sRPE, RPE and its derivatives have been used evenly from 2010 to 2022. However, HR was mostly used (in six out of seven studies) from 2019 to 2021. See Figure 2.

Athletes' readiness

Analyses of data collected from wellness or well-being questionnaires (WB) (Hooper & Mackinnon, 1995; McLean, Coutts, Kelly, McGuigan, & Cormack, 2010) were the most observed methods for the assessment of athletes' readiness in 14 studies (Brandão, et al., 2019; Clemente, et al., 2019, 2020; de Leeuw, et al., 2022; Duarte, Coimbra, et al., 2019; Edmonds, et al., 2021; Haraldsdottir, et al., 2021; Kupperman, et al., 2021; Mendes, et al., 2018; Rabbani, et al., 2021; Roy, et al., 2019; Tavares, et al., 2018; Timoteo, et al., 2017; Ungureanu, Lupo, et al., 2021), followed by the Total Quality Recovery Scale (TQR), used in 10 studies (Andrade, et al., 2021; Brandão, et al., 2019; Cardoso, et al., 2021; Debien, et al., 2018; Duarte, Coimbra, et al., 2019; Freitas, et al., 2014; Horta, et al., 2020; Lacerda, et al., 2015; Timoteo, et al., 2017, 2021). Other questionnaires such as the Recovery Stress Questionnaire for Sports (REST-Q-sports) were used in four studies (Freitas, et al., 2014; Freitas, Nakamura, et al., 2015; Horta, Bara Filho, Coimbra, Miranda, et al., 2019; van der Does, et al., 2017), the Profile

of Mood States (POMS) in two studies (Aoki, et al., 2017; Horta, et al., 2020) and lastly the Daily Analyses of Life Demands of Athletes (DALDA) in one study (Moreira, et al., 2010). Other different scales were also found in our review, with one study each: the Rating of Perceived Recovery (RPR) (Kraft, et al., 2020), the Visual Analogue Scale (VAS) for mental fatigue (Coyne, et al., 2021) and the Perceived Recovery State (PRS) (Cardoso, et al., 2021). Objective measures were also collected, via heart rate variability (HRV) (Cardoso, et al., 2021; Edmonds, et al., 2021; Hernández-Cruz, et al., 2017; Rabbani, et al., 2021) and countermovement jump (CMJ) (Carroll, et al., 2019; Freitas, et al., 2014; Freitas, Nakamura, et al., 2015; Rabbani, et al., 2021; Tavares, et al., 2018) in four and five studies, respectively. See Table 3.

If a chronological order of use in studies is implemented for readiness measures, the first method detected is DALDA in a 2010 study, followed by the REST-Q sport, TQR and CMJ from 2014 to 2019-2021 and lastly WB and HRV from 2017 to 2021. See Figure 2.

External load

The most studied variables to monitor external workload were vertical displacement variables, specifically, jump count and/or jump load. In 12 studies microensors were used to measure workload (predominantly VERT Classic and Catapult Sports' Optimeye 5S) (Herring & Fukuda, 2022; Kupperman, et al., 2021; Libs, et al., 2019; Lima, et al., 2019, 2021, Lima et al., 2020; Piatti, et al., 2021; Rabello, et al., 2019; Sanders, et al., 2018; Skazalski, et al., 2018; Taylor, et al., 2022; Vlantes & Readdy, 2017), but also: jump height, establishing height thresholds, detect jump type and measure jump intensity from the devices' integrated gyroscope, magnetometer and tri-axial accelerometer. Other metrics such as player load, vertical accelerations, high impacts, high impacts % and explosive efforts could also be obtained from their software.

Video analysis was also used in seven studies (Bahr & Bahr, 2014; Garcia-de-Alcaraz, et al., 2020; Lima, et al., 2021; Rabello, et al., 2019; Taylor, et al., 2019; Ungureanu, Lupo, et al., 2021; Wolfe, et al., 2019) for jump count/load, the detection of the type of jumps/landings, jumps by position, the calculation of distances covered by players, technical actions quantification (sets, spikes, serves, blocks, digs, receptions, defences). Other methods were also observed for external workload monitoring, including swing count (Wolfe, et al., 2019), Changes of direction, accelerations, decelerations, and high-intensity efforts (Kupperman, et al., 2021), data volley variables (defences, receptions, digs...) (Lima, et al., 2021) and sets, repetitions and loads in the gym and/or court sessions (de Leeuw, et al.,

2022). See Table 3.

Finally, if we look at the evolution of external methods through the years, the first method to be observed in volleyball studies on external load was the video analysis of jump load in 2014. In 2017 microensors started to appear, and from there, all the methods are evenly used from 2019 onwards. See Figure 2.

Discussion and conclusions

This systematic review seeks to address the lack of consensus on the selection of training and competition load monitoring tools (Fox, Stanton, Sargent, Wintour, & Scanlan, 2018) as well as on the methods for assessing the readiness of volleyball players. The findings of this review are intended to provide valuable information for sports professionals to facilitate their informed decision making about their training plans (Jeffries, et al., 2022). To achieve this purpose, we presented the most commonly used methods found in scientific literature and their trend of use over time with the intention to provide an updated record of tools that can be employed by any volleyball team.

In this review, three clearly defined types of tools have been identified and described in the current literature (Jeffries, et al., 2022). These tools are divided into those that monitor internal loads, those that focus on external loads, and those used to assess players' readiness.

In volleyball, it is common to use multiple monitoring tools (Clemente, et al., 2019; Mendes, et al., 2018). To make more accurate planning decisions, it is recommended to combine tools that measure internal loads, external loads, and players' readiness (Burgess, 2017; De Beéck, et al., 2019; Fox, et al., 2018; S. Ryan, Kempton, & Coutts, 2021; Saw, Main, & Gastin, 2016).

The sRPE is a popular tool for internal monitoring due to its simplicity and ability to provide detailed information. For assessing players' readiness, the TQR scale, wellness questionnaires and the CMJ are prominent options. For external loads, it is important to measure a variety of actions, ideally using microensors in all three axes of motion.

Objective measurement, through technological advances in devices, shows a steady increase in the literature since 2017. Especially in technology focused on quantifying workloads during on-court sessions. Among the most prominent contributions of technology is the ability to provide real-time information (Garcia-de-Alcaraz, et al., 2020; Lima, et al., 2021; Ungureanu, Lupo, et al., 2021) and to quantify load in all spatial axes (Kupperman, et al., 2021).

However, subjective methods through questionnaires have continued to be used. In some cases, there is an explicit confrontation between the objective and subjective sources. For example, although

objective methods exist to measure internal load, such as heart rate for the calculation of TRIMP (Bara Filho, et al., 2013; Duarte, Alves, et al., 2019; González, et al., 2005; Kraft, et al., 2020; Libs, et al., 2019; Rodríguez-Marroyo, et al., 2014), a preference for the use of sRPE has been observed.

Technological advances in the search for greater objectivity in measurement are intrinsic to scientific research and sports training.

The use of subjective tools such as the sRPE and wellness questionnaires has also been found to affect self-awareness and, in addition, to promote the development of self-regulation (Vavassori, Moreno, & Ureña, 2023).

Therefore, from a perspective based on subjective insight, there is a phenomenological approach (Sousa, 2014; Vavassori, et al., 2023; Zahavi, 2020) that could provide value in terms of self-regulation.

Studies on self-regulation have highlighted its relevance in sports development, performance and readiness (Balk & Englert, 2020; Harrison, et al., 2022). In addition, its importance has been evidenced in issues related to well-being (Crawford, Tripp, Gierc, & Scott, 2021), which includes the aspects assessed in the wellness questionnaires analyzed in this review. However, it was not that there was hidden knowledge about the value of self-regulation. Rather, there was a comfort and/or accessibility that was not refuted by the technology. Hence the importance of giving added value to qualitative instruments.

Possibly, the extensive use of sRPE in volleyball (Pisa, et al., 2022) may also be due to the existing relationship between various tools regardless of their objective or subjective nature.

Although the focus in volleyball has been on quantifying jumps for years, it is relevant to note that less than 50% of a players' total load on the court comes from jumps, as significant load occurs during horizontal movements (Vlantes & Readdy, 2017). Volleyball is characterized by a series of small movements, accelerations, decelerations and changes of direction that generate high stress on players, and thanks to technological advances, these can be detected through microsensors (Kupperman, et al., 2021).

On the other hand, obtaining real-time information allows for faster and highly individualized training decisions. Individualization in the monitoring and planning of sessions is crucial, since, for example, the volume and intensity of jumps vary significantly depending on the role of each player (Skazalski, et al., 2018; Vlantes & Readdy, 2017). This highlights the importance of establishing player-specific load thresholds (Brito, Hertzog & Nassis, 2016; Kellmann, et al., 2018). However, we should not underestimate another potential benefit of the immediate feedback offered by some tech-

nologies, such as the stimulation of self-motivation. Motivation theories distinguish between mastery-focused motivation and ego-focused motivation (Ryan & Deci, 2000). In the case of volleyball student-athletes using objective tools, it has been observed that their motivation is mainly focused on outperforming their teammates (Vavassori, et al., 2023). Therefore, we should not dismiss the motivational contributions they can derive from technology and objective methods.

Although these two perspectives (objective and subjective) are interconnected, as information from the objective world can influence human consciousness and decisions, technological advances transform the subjective into objective information for information systems (Xu, et al., 2023). Therefore, although the relationship between these two dimensions is complex, the information is simultaneously subjective and objective (Bates, 2006).

In summary, using the sRPE (and its derivatives), TQR/WB, CMJ would create a monitoring system sufficient for teams in developmental stages and at a reduced cost. Furthermore, it is worth insisting on taking advantage of the added value in terms of self-regulation and motivation provided by the use of qualitative instruments. However, the combination of these tools with microsensors would result in a complete and real-time monitoring system for decision-making of the volleyball team staff.

Limitations and strengths

Because of the reduced number of researches regarding monitoring in the sport of volleyball compared to other team sports (soccer, rugby, Australian football), limitations may arise in the current review. Many studies identified used a limited number of participants. Although volleyball teams usually have 12-14 players, and collecting data from more than one team might not be feasible, small sample conclusions should be taken with caution. Also, there might have been some selection bias as in team sports, the composition of the teams is already set and players are not selected randomly. Another possible selection bias could have arisen from the decision to use only one author for the initial selection of studies. Even though PRISMA allows the use of a single author for this stage, some studies might have been wrongly included or excluded during the process.

Comparison between studies is not advised, as findings in studies with different statistical analyses may be complicated. Meta-analysis is suggested in the future to solve this issue. However, to minimize this effect, article quality was assessed to reduce bias and include higher standard research in the results. We may have incurred in risk of bias by not executing a dual and independent screening.

Nevertheless, we are confident that the conclusions of this review have not been affected by these methodological limitations.

Important to notice that, due to different playing levels, ages and genders, training control methods should be adapted to each team, situation, level, goals, and limiting factors.

Despite these constraints, we consider the information provided in this systematic review may contribute to increasing team performance, avoiding non-functional overreaching and hence, mitigating injury occurrence by selecting monitoring methods supported by science and used by professionals in elite and development teams in volleyball. It may also help coaches in selecting the best available method to monitor the load and readiness of their teams.

Future directions

In this review, two studies quantified load on 3 axes measuring vertical and horizontal displacement with wearable microsensors, concluding vertical displacement loads cover less than half of the training and match loads. Consequently, horizontal movements create greater workloads (Kupperman, et al., 2021; Vlantes & Readdy, 2017).

Thus, more research is needed as most studies in the past focused solely on jumps.

All sessions of a training week (gym, individual training, rehab) either in the preseason or competitive period should be investigated further. Since the current research has limited the information to on-court sessions only, we feel an immense quantity of load is discarded. Furthermore, some studies did not include match loads in the weekly load calculation, creating false load information. Keep in mind that match load usually is the highest load of the week (Brito, et al., 2016; Fessi & Moalla, 2018; Murphy, Duffield, Kellett, & Reid, 2016).

From our search, there was no study observing training loads and readiness distribution during the off-season. Collecting information during this period might assist in anticipating the planning of workloads for the preseason, normally the period with the highest volume of the year (Andrade, et al., 2021; Aoki, et al., 2017; Horta, Bara Filho, Coimbra, Miranda, et al., 2019). By doing so, coaches should be able to avoid excessive spikes in load by having access to workload data from the transition period. Studies in the future may shed light on what methods are used to monitor players when are away from team settings.

References

- Andrade, D.M., Fernandes, G., Miranda, R., Reis Coimbra, D., & Bara Filho, M.G. (2021). Training load and recovery in volleyball during a competitive season. *Journal of Strength and Conditioning Research*, 35(4), 1082-1088. <https://doi.org/10.1519/JSC.0000000000002837>
- Aoki, M.S., Arruda, A.F.S., Freitas, C.G., Miloski, B.D., Marcelino, P.R., Drago, G., Drago, M., & Moreira, A. (2017). Monitoring training loads, mood states, and jump performance over two periodized training mesocycles in elite young volleyball players. *International Journal of Sports Science and Coaching*, 12(1), 130-137. <https://doi.org/10.1177/1747954116684394>
- Bahr, M.A., & Bahr, R. (2014). Jump frequency may contribute to risk of jumper's knee: A study of interindividual and sex differences in a total of 11 943 jumps video recorded during training and matches in young elite volleyball players. *British Journal of Sports Medicine*, 48(17), 1322-1326. <https://doi.org/10.1136/bjsports-2014-093593>
- Balk, Y.A., & Englert, C. (2020). Recovery self-regulation in sport: Theory, research, and practice. *International Journal of Sports Science and Coaching*, 15(2), 273-281. <https://doi.org/10.1177/1747954119897528>
- Bara Filho, M.G., de Andrade, F.C., Nogueira, R.A., & Nakamura, F.Y. (2013). Comparison of different methods of internal load control in volleyball players. *Revista Brasileira de Medicina do Esporte*, 19(2), 143-146. <https://doi.org/10.1590/S1517-86922013000200015>
- Bates, M.J. (2006). Fundamental forms of information. *Journal of the American Society for Information Science and Technology*, 57(8), 1033-1045. <https://doi.org/10.1002/asi.20369>
- Bourdon, P.C., Cardinale, M., Murray, A., Gatin, P.B., Kellmann, M., Varley, M.C., Gabbett, T., Coutts, A.J., Burgess, D.J., Gregson, W., & Cable, N.T. (2017). Monitoring athlete training loads: Consensus statement. *International Journal of Sports Physiology and Performance*, 12(Supplement 2), 161-170. <https://doi.org/10.1123/IJSP.2017-0208>
- Brandão, F.M., Da Cunha, V.F., Timoteo, T.F., Duarte, T.S., Dias, B.M., Coimbra, D.R., Miranda, R., & Bara Filho, M.G. (2019). Comportamento da carga de treinamento, recuperação e bem-estar em atletas profissionais de voleibol em semanas com e sem jogos. [Behavior of the training load, recovery and well-being in volleyball professional athletes in weeks with and without matches]. *Educación Física y Ciencia*, 20(4), e063. <https://doi.org/10.24215/23142561e063>

- Brito, J., Hertzog, M., & Nassis, G.P. (2016). Do match-related contextual variables influence training load in highly trained soccer players? *Journal of Strength and Conditioning Research*, 30(2), 393-399. <https://doi.org/10.1519/JSC.0000000000001113>
- Burgess, D.J. (2017). The research doesn't always apply: Practical solutions to evidence-based training-load monitoring in elite team sports. *International Journal of Sports Physiology and Performance*, 12(Supplement 2), 136-141. <https://doi.org/10.1123/ijssp.2016-0608>
- Cardoso, A.S., Berriel, G.P., Schons, P., Costa, R.R., & Krueel, L.F.M. (2021). Recovery behavior after matches for returning to training in volleyball athletes. *Archivos de Medicina Del Deporte*, 38(5), 343-349. <https://doi.org/10.18176/ARCHMEDDEPORTE.00061>
- Carroll, K.M., Wagle, J.P., Sole, C.J., & Stone, M.H. (2019). Intrasession and intersession reliability of countermovement jump testing in division-I volleyball athletes. *Journal of Strength and Conditioning Research*, 33(11), 2932-2935. <https://doi.org/10.1519/JSC.0000000000003353>
- Castellano, J., Alvarez-Pastor, D., & Bradley, P.S. (2014). Evaluation of research using computerised tracking systems (amisco® and prozone®) to analyse physical performance in elite soccer: A systematic review. *Sports Medicine*, 44(5), 701-712. <https://doi.org/10.1007/s40279-014-0144-3>
- Castello, M., Reed, J.P., Lund, R., & Mack, M. (2018). Relationship between physical training, ratings of perceived exertion, and mental toughness in female NCAA Division I volleyball players. *Sport Journal*, 21, 1-8.
- Çelebi, M.M., & Aksu, A. (2018). Incidence of injuries in female and male volleyball players—Prospective preliminary study. / Kadın ve Erkek Voleybolcularda Yaralanma İnsidansı—Prospektif Ön Çalışma. *Sport Hekimligi Dergisi/Turkish Journal of Sports Medicine*, 53(3), 109-115.
- Clemente, F.M., Mendes, B., Nikolaidis, P.T., Calvete, F., Carriço, S., & Owen, A.L. (2017). Internal training load and its longitudinal relationship with seasonal player wellness in elite professional soccer. *Physiology and Behavior*, 179, 262-267. <https://doi.org/10.1016/j.physbeh.2017.06.021>
- Clemente, F.M., Mendes, B., Palao, J.M., Silverio, A., Carriço, S., Calvete, F., & Nakamura, F.Y. (2019). Seasonal player wellness and its longitudinal association with internal training load: Study in elite volleyball. *Journal of Sports Medicine and Physical Fitness*, 59(3), 345-351. <https://doi.org/10.23736/S0022-4707.18.08312-3>
- Clemente, F.M., Silva, A.F., Clark, C.C.T., Conte, D., Ribeiro, J., Mendes, B., & Lima, R.F. (2020). Analyzing the seasonal changes and relationships in training load and wellness in elite volleyball players. *International Journal of Sports Physiology and Performance*, 15(5), 731-740. <https://doi.org/10.1123/ijssp.2019-0251>
- Coyne, J.O.C., Coutts, A.J., Newton, R.U., & Haff, G.G. (2021). The influence of mental fatigue on sessional ratings of perceived exertion in elite open and closed skill sports athletes. *Journal of Strength and Conditioning Research*, 35(4), 963-969. <https://doi.org/10.1519/JSC.0000000000003980>
- Crawford, A., Tripp, D.A., Gierc, M., & Scott, S. (2021). The influence of mental toughness and self-regulation on post-season perceptions in varsity athletes. *Journal of American College Health*, 71(4), 1036-1044. <https://doi.org/10.1080/07448481.2021.1920596>
- de Andrade, F.C., Alves Nogueira, R., Coimbra, D.R., Miloski, B.D., Freitas, V.H. de, & Bara Filho, M.G. (2014). Internal training load: Perception of volleyball coaches and athletes. / Carga interna de treinamento: percepção de técnicos e atletas de voleibol. *Brazilian Journal of Kineanthropometry and Human Performance*, 16(6), 638-647. <https://doi.org/10.5007/1980-0037.2014v16n6p638>
- De Beéck, T.O., Jaspers, A., Brink, M.S., Frencken, W., Staes, F., Davis, J.J., & Helsen, W.F. (2019). Predicting future perceived wellness in professional soccer: The role of preceding load and wellness. *International Journal of Sports Physiology and Performance*, 14(8), 1074-1080. <https://doi.org/10.1123/ijssp.2017-0864>
- Debien, P.B., Mancini, M., Coimbra, D.R., Freitas, D.G.S., Miranda, R., & Bara Filho, M.G. (2018). Monitoring training load, recovery, and performance of Brazilian professional volleyball players during a season. *International Journal of Sports Physiology and Performance*, 13(9), 1182-1189. <https://doi.org/10.1123/ijssp.2017-0504>
- de Leeuw, A.-W., van der Zwaard, S., van Baar, R., & Knobbe, A. (2022). Personalized machine learning approach to injury monitoring in elite volleyball players. *European Journal of Sport Science*, 22(4), 511-520. <https://doi.org/10.1080/17461391.2021.1887369>
- Duarte, T.S., Alves, D.L., Coimbra, D.R., Miloski, B.D., Bouzas Marins, J. C., & Bara Filho, M.G. (2019). Technical and tactical training load in professional volleyball players. *International Journal of Sports Physiology and Performance*, 14(10), 1338-1343. <https://doi.org/10.1123/ijssp.2019-0004>
- Duarte, T.S., Coimbra, D.R., Miranda, R., Toledo, H.C., Werneck, F.Z., Freitas, D.G.S., & Bara Filho, M.G. (2019). Monitoring training load and recovery in volleyball players during a season. [Monitoramento da carga de treinamento e recuperação em jogadores de voleibol durante uma temporada.] *Revista Brasileira de Medicina do Esporte*, 25(3), 226-229. <https://doi.org/10.1590/1517-869220192503195048>
- Edmonds, R., Schmidt, B., & Siedlik, J. (2021). Eligibility classification as a factor in understanding student-athlete responses to collegiate volleyball competition. *Sports (Basel, Switzerland)*, 9(3), 43. <https://doi.org/10.3390/sports9030043>
- Eliakim, A., Portal, S., Zadik, Z., Rabinowitz, J., Adler-Portal, D., Cooper, D.M., Zaldivar, F., & Nemet, D. (2009). The effect of a volleyball practice on anabolic hormones and inflammatory markers in elite male and female adolescent players. *Journal of Strength and Conditioning Research*, 23(5), 1553-1559. <https://doi.org/10.1519/JSC.0b013e3181aa1bcb>

- Fessi, M.S., & Moalla, W. (2018). Postmatch perceived exertion, feeling, and wellness in professional soccer players. *International Journal of Sports Physiology and Performance*, 13(5), 631-637. <https://doi.org/10.1123/ijsp.2017-0725>
- Fessi, M.S., Nouira, S., Dellal, A., Owen, A.L., Elloumi, M., & Moalla, W. (2016). Changes of the psychophysical state and feeling of wellness of professional soccer players during pre-season and in-season periods. *Research in Sports Medicine*, 24(4), 375-386. <https://doi.org/10.1080/15438627.2016.1222278>
- Foster, C., Florhaug, J.A., Franklin, J., Gottschall, L., Hrovatin, L.A., Parker, S., Doleshal, P., & Dodge, C. (2001). A new approach to monitoring exercise training. *Journal of Strength and Conditioning Research*, 15(1), 109-115. <https://doi.org/10.1519/00124278-200102000-00019>
- Fox, J.L., Stanton, R., Sargent, C., Wintour, S.A., & Scanlan, A.T. (2018). The association between training load and performance in team sports: A systematic review. *Sports Medicine*, 48(12), 2743-2774. <https://doi.org/10.1007/s40279-018-0982-5>
- Freitas, V.H. de, Miloski, B.D., & Bara Filho, M.G. (2015). Monitoramento da carga interna de um período de treinamento em jogadores de voleibol. [Internal training load monitoring across a period training in volleyball players]. *Revista Brasileira de Educação Física e Esporte*, 29(1), 5-12. <https://doi.org/10.1590/1807-55092015000100005>
- Freitas, V.H. de, Nakamura, F.Y., de Andrade, F.C., Pereira, L.A., Coimbra, D.R., & Bara Filho, M.G. (2015). Pre-competitive physical training and markers of performance, stress and recovery in young volleyball athletes. / Treinamento físico pré-competitivo e marcadores de desempenho, estresse e recuperação em jovens atletas de voleibol. *Revista Brasileira de Cineantropometria e Desempenho Humano*, 17(1), 31-40. <https://doi.org/10.5007/1980-0037.2015v17n1p31>
- Freitas, V.H. de, Nakamura, F.Y., Miloski, B.D., Samulski, D., & Bara Filho, M.G. (2014). Sensitivity of physiological and psychological markers to training load intensification in volleyball players. *Journal of Sports Science and Medicine*, 13(3), 571-579.
- García-de-Alcaraz, A., Ramírez-Campillo, R., Rivera-Rodríguez, M., & Romero-Moraleda, B. (2020). Analysis of jump load during a volleyball season in terms of player role. *Journal of Science and Medicine in Sport*, 23(10), 973-978. <https://doi.org/10.1016/j.jsams.2020.03.002>
- García-de-Alcaraz, A., Valadés, D., & Palao, J.M. (2017). Evolution of game demands from young to elite players in men's volleyball. *International Journal of Sports Physiology and Performance*, 12(6), 788-795. <https://doi.org/10.1123/ijsp.2016-0027>
- Gielen, J., Mehuys, E., Berckmans, D., Meeusen, R., & Aerts, J.-M. (2022). Monitoring internal and external load during volleyball competition. *International Journal of Sports Physiology and Performance*, 17(4), 640-645. <https://doi.org/10.1123/ijsp.2021-0217>
- González, C., Ureña, A., Llop, F., García, J.M., Martín, A., & Navarro, F. (2005). Physiological characteristics of libero and central volleyball players. *Biology of Sport*, 22(1), 13-27.
- Hamlin, M.J., Wilkes, D., Elliot, C.A., Lizamore, C.A., & Kathiravel, Y. (2019). Monitoring training loads and perceived stress in young elite university athletes. *Frontiers in Physiology*, 10, 34. <https://doi.org/10.3389/fphys.2019.00034>
- Hank, M., Zahalka, F., & Maly, T. (2015). Comparison of spikers' distance covered in elite female volleyball. [Usporedba pokrivene udaljenosti 'spikera' u elitnoj ženskoj odbojci.]. *Sport Science*, 8, 102-106.
- Háp, P., Stejskal, P., Jakubec, A., Hap, P., Stejskal, P., & Jakubec, A. (2011). Volleyball players training intensity monitoring through the use of spectral analysis of heart rate variability during a training microcycle. *Acta Gymnica*, 41(3), 33-38. <https://doi.org/10.5507/ag.2011.018>
- Haraldsdottir, K., Sanfilippo, J., McKay, L., & Watson, A.M. (2021). Decreased sleep and subjective well-being as independent predictors of injury in female collegiate volleyball players. *Orthopaedic Journal of Sports Medicine*, 9(9). <https://doi.org/10.1177/232596712111029285>
- Harrison, C., Ruddock-Hudson, M., Mayes, S., O'Halloran, P., Ferrar, K., Ruddock, S., & Cook, J. (2022). An exploration of the perceptions and experiences of professional ballet dancers using a wellness monitoring application. *Qualitative Research in Sport, Exercise and Health*, 14(7), 1196-1212. <https://doi.org/10.1080/2159676X.2022.2111456>
- Hernández-Cruz, G., Quezada-Chacón, J.T., González-Fimbres, R.A., Flores-Miranda, F.J., Naranjo-Orellana, J., & Rangel-Colmenero, B.R. (2017). Effect of consecutive matches on heart rate variability in elite volleyball players. *Revista de Psicoogía Del Deporte*, 26(2), 9-14.
- Herring, C.H., & Fukuda, D.H. (2022). Monitoring competition jump load in division I female collegiate volleyball athletes. *Journal of Science in Sport and Exercise*, 4, 221-230. <https://doi.org/10.1007/s42978-021-00152-y>
- Hooper, S.L., & Mackinnon, L.T. (1995). Monitoring overtraining in athletes: Recommendations. *Sports Medicine*, 20(5), 321-327. <https://doi.org/10.2165/00007256-199520050-00003>
- Horta, T.A.G., Bara Filho, M.G., Coimbra, D.R., Miranda, R., & Werneck, F.Z. (2019). Training load, physical performance, biochemical markers, and psychological stress during a short preparatory period in Brazilian elite male volleyball players. *Journal of Strength and Conditioning Research*, 33(12), 3392-3399. <https://doi.org/10.1519/JSC.0000000000002404>

- Horta, T.A.G., Bara Filho, M.G., Coimbra, D.R., Werneck, F.Z., & Miranda, R. (2019). Training load profile in high performance volleyball: A case study. [Perfil da carga de treinamento no voleibol de alto rendimento: um estudo de caso] [Perfil de la carga de entrenamiento en el voleibol de alto rendimiento: un estudio de caso.]. *Revista Brasileira de Ciencias do Esporte*, 41(4), 419-426. <https://doi.org/10.1016/j.rbce.2018.06.008>
- Horta, T.A.G., Bara Filho, M.G., Miranda, R., Coimbra, D.R., & Werneck, F.Z. (2017). Influence of vertical jump in the perception of the internal volleyball training load. [Influência dos saltos verticais na percepção da carga interna de treinamento no voleibol.] *Revista Brasileira de Medicina do Esporte*, 23(5), 403-406. <https://doi.org/10.1590/1517-869220172305172132>
- Horta, T.A.G., Coimbra, D.R., Miranda, R., Werneck, F.Z., & Bara Filho, M.G. (2017). Is the internal training load different between starters and nonstarters volleyball players submitted to the same external load training? A case study. / A carga interna de treinamento é diferente entre atletas de voleibol titulares e reservas? Um estudo. *Brazilian Journal of Kineanthropometry and Human Performance*, 19(4), 395-405. <https://doi.org/10.5007/1980-0037.2017v19n4p395>
- Horta, T.A.G., de Lima, P.H.P., Matta, G.G., de Freitas, J.V., Miloski, B.D., Vianna, J.M., Toledo, H.C., Miranda, R., Timoteo, T.F., & Bara Filho, M.G. (2020). Training load impact on recovery status in professional volleyball athletes. *Revista Brasileira de Medicina do Esporte*, 26(2), 158-161. <https://doi.org/10.1590/1517-869220202602209364>
- Hurd, W., Hunter-Giordano, A., Axe, M., & Snyder-Mackler, L. (2009). Data-based interval hitting program for female college volleyball players. *Sports Health*, 1(6), 522-530. <https://doi.org/10.1177/1941738109351171>
- Jeffries, A.C., Marcora, S.M., Coutts, A.J., Wallace, L., McCall, A., & Impellizzeri, F.M. (2022). Development of a revised conceptual framework of physical training for use in research and practice. *Sports Medicine*, 52(4), 709-724. <https://doi.org/10.1007/s40279-021-01551-5>
- Jeffries, A.C., Wallace, L., Coutts, A.J., McLaren, S.J., McCall, A., & Impellizzeri, F.M. (2020). Athlete-reported outcome measures for monitoring training responses: A systematic review of risk of bias and measurement property quality according to the COSMIN guidelines. *International Journal of Sports Physiology and Performance*, 15(9), 1203-1215. <https://doi.org/10.1123/IJSPP.2020-0386>
- Kellmann, M., Bertollo, M., Bosquet, L., Brink, M., Coutts, A.J., Duffield, R., ..., & Beckmann, J. (2018). Recovery and performance in sport: Consensus statement. *International Journal of Sports Physiology and Performance*, 13(2), 240-245. <https://doi.org/10.1123/ijssp.2017-0759>
- Kraft, J.A., Laurent, M.C., Green, J.M., Helm, J., Roberts, C., & Holt, S. (2020). Examination of coach and player perceptions of recovery and exertion. *Journal of Strength and Conditioning Research*, 34(5), 1383-1391. <https://doi.org/10.1519/JSC.0000000000002538>
- Kupperman, N., Curtis, M.A., Saliba, S.A., & Hertel, J. (2021). Quantification of workload and wellness measures in a women's collegiate volleyball season. *Frontiers in Sports and Active Living*, 3, 702419. <https://doi.org/10.3389/fspor.2021.702419>
- Lacerda, R.P., Duarte, T.S., Coimbra, D.R., Timoteo, T.F., Marins, J.C.B., Miranda, R., & Bara Filho, M.G. (2015). Comportamento da recuperação de atletas profissionais de voleibol em semanas com jogos e sem jogos. [Behavior of recovery in volleyball professional athletes in weeks with and without matches]. *Coleção Pesquisa Em Educação Física*, 14(2), 23-30.
- Libs, H., Boos, B., Shipley, F., Peacock, C.A., & Sanders, G.J. (2019). Variability in Preseason jump loads and heart rate intensities in Division I volleyball. *Journal of Exercise and Nutrition*, 2(2), 8.
- Lima, R., Castro, H. de O., Afonso, J., Costa, G.D.C.T., Matos, S., Fernandes, S., & Clemente, F.M. (2021). Effects of congested fixture on men's volleyball load demands: Interactions with sets played. *Journal of Functional Morphology and Kinesiology*, 6(2), 53. <https://doi.org/10.3390/jfmk6020053>
- Lima, R.F., Palao, J.M.J., Castro, H., & Clemente, F.M. (2019). Measuring the training external jump load of elite male volleyball players: An exploratory study in Portuguese League. / Medición de la carga externa de entrenamiento de los jugadores de voleibol masculino de élite: un estudio exploratorio en la Liga Port. *Retos: Nuevas Perspectivas de Educación Física, Deporte y Recreación*, 36(36), 454-458.
- Lima, R.F., Silva, A.F., Afonso, J., Castro, H., & Clemente, F.M. (2020). External and internal load and their effects on professional volleyball training. *International Journal of Sports Medicine*, 41(7), 468-474. <https://doi.org/10.1055/a-1087-2183>
- Maksimenko, I.G., Maksimenko, G.N., Zhilina, L.V., & Bayeva, D.N. (2019). Training workload controls applicable in elite women's volleyball. *Teoriya i Praktika Fizicheskoy Kultury*, 7, 74-76.
- Malisoux, L., Frisch, A., Urhausen, A., Seil, R., & Theisen, D. (2013). Monitoring of sport participation and injury risk in young athletes. *Journal of Science and Medicine in Sport*, 16(6), 504-508. <https://doi.org/10.1016/j.jsams.2013.01.008>
- McLean, B.D., Coutts, A.J., Kelly, V., McGuigan, M.R., & Cormack, S.J. (2010). Neuromuscular, endocrine, and perceptual fatigue responses during different length between-match microcycles in professional rugby league players. *International Journal of Sports Physiology and Performance*, 5(3), 367-383. <https://doi.org/10.1123/ijssp.5.3.367>

- Mendes, B., Palao, J.M., Silverio, A., Owen, A.L., Carriço, S., Calvete, F., & Clemente, F.M. (2018). Daily and weekly training load and wellness status in preparatory, regular and congested weeks: A season-long study in elite volleyball players. *Research in Sports Medicine*, 26(1), 462-473. <https://doi.org/10.1080/15438627.2018.1492393>
- Moreira, A., de Freitas, C.G., Nakamura, F.Y., & Aoki, M.S. (2010). Percepção de esforço da sessão e a tolerância ao estresse em jovens atletas de voleibol e basquetebol. [Session RPE and stress tolerance in young volleyball and basketball players]. *Revista Brasileira de Cineantropometria e Desempenho Humano*, 12(5), 345-351. <https://doi.org/10.5007/1980-0037.2010v12n5p345>
- Moreira, A., Freitas, C.G., Nakamura, F.Y., Drago, G., Drago, M., & Aoki, M.S. (2013). Effect of match importance on salivary cortisol and immunoglobulin a response in elite young volleyball players. *Journal of Strength and Conditioning Research*, 27(1), 202-207. <https://doi.org/10.1519/JSC.0b013e31825183d9>
- Mortatti, A.L., Pinto, J.C.B.L., Lambertucci, R., Hirabara, S.M., & Moreira, A. (2018). Does a congested fixture schedule affect psychophysiological parameters in elite volleyball players? [Est-ce qu'un calendrier surcharge affecte les paramètres psychophysologiques chez les joueurs d'élite de volleyball?] *Science and Sports*, 33(4), 258-264. <https://doi.org/10.1016/j.scispo.2018.02.015>
- Mroczek, D., Januszkiewicz, A., Kawczynski, A.S., Borysiuk, Z., Chmura, J., Kawczynski, A.S., Borysiuk, Z., & Chmura, J. (2014). Analysis of male volleyball players' motor activities during a top level match. *Journal of Strength and Conditioning Research*, 28(8), 2297-2305. <https://doi.org/10.1519/JSC.0000000000000425>
- Murphy, A.P., Duffield, R., Kellett, A., & Reid, M. (2016). A comparison of the perceptual and technical demands of tennis training, simulated match play, and competitive tournaments. *International Journal of Sports Physiology and Performance*, 11(1), 40-47. <https://doi.org/10.1123/ijsp.2014-0464>
- Noce, F., Dos Santos, I.C., Samulski, D., De Carvalho, S.L.F., Dos Santos, R.V.T., & De Mello, M.T. (2008). Monitoring levels of stress and overtraining in an elite Brazilian female volleyball athlete: Case study. *Revista de Psicologia Del Deporte*, 17(1), 25-41.
- Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., ..., & Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*, 372, n71. <https://doi.org/10.1136/bmj.n71>
- Piatti, M., Ambrosi, E., Dedda, G., Omeljaniuk, R.J., Turati, M., Bigoni, M., & Gaddi, D. (2021). Jump performance during a season in elite volleyball players. *The Journal of Sports Medicine and Physical Fitness*, 62(5), 602-608. <https://doi.org/10.23736/S0022-4707.21.12268-6>
- Pires, D.A., Bara Filho, M.G., Debien, P.B., Coimbra, D.R., & Ugrinowitsch, H. (2016). Burnout and coping among volleyball athletes: A longitudinal analysis. [Burnout e coping em atletas de voleibol: Uma análise longitudinal.] *Revista Brasileira de Medicina do Esporte*, 22(4), 277-281. <https://doi.org/10.1590/1517-869220162204158756>
- Pires, D.A., & Ugrinowitsch, H. (2021). Burnout and coping perceptions of volleyball players throughout an annual sport season. *Journal of Human Kinetics*, 79(1), 249-257. <https://doi.org/10.2478/hukin-2021-0078>
- Pisa, M.F., Zecchin, A.M., Gomes, L.G., Norberto, M.S., & Puggina, E.F. (2022). Internal load in male professional volleyball: A systematic review. *The Journal of Sports Medicine and Physical Fitness*, 62(11), 1465-1473. <https://doi.org/10.23736/s0022-4707.22.12885-9>
- Podstawski, R., Boraczynski, M., Nowosielska-Swadzba, D., & Zwolinska, D. (2014). Heart rate variability during pre-competition and competition periods in volleyball players. *Biomedical Human Kinetics*, 6(1), 19-26. <https://doi.org/10.2478/bhk-2014-0004>
- Rabbani, M., Agha-Alinejad, H., Gharakhanlou, R., Rabbani, A., & Flatt, A.A. (2021). Monitoring training in women's volleyball: Supine or seated heart rate variability? *Physiology and Behavior*, 240, 113537. <https://doi.org/10.1016/j.physbeh.2021.113537>
- Rabello, L.M., Zwerver, J., Stewart, R.E., van den Akker-Scheek, I., & Brink, M.S. (2019). Patellar tendon structure responds to load over a 7-week preseason in elite male volleyball players. *Scandinavian Journal of Medicine and Science in Sports*, 29(7), 992-999. <https://doi.org/10.1111/sms.13428>
- Reynoso-Sánchez, L.-F., Hernández-Cruz, G., López-Walle, J., Rangel-Colmenero, B., Quezada-Chacón, J.-T., & Jaenes Sánchez, J.C. (2016). Balance de estrés-recuperación en jugadores universitarios de voleibol durante una temporada. [Recovery-stress balance throughout a season in volleyball university players.] *Retos*, 30(30), 193-197. <https://doi.org/10.47197/retos.v0i30.50244>
- Rodríguez-Marroyo, J.A., Medina, J., García-López, J., García-Tormo, J.V., & Foster, C. (2014). Correspondence between training load executed by volleyball players and the one observed by coaches. *Journal of Strength and Conditioning Research*, 28(6), 1588-1594. <https://doi.org/10.1519/JSC.0000000000000324>
- Roy, X., Caya, O., Charron, J., Comtois, A.S., & Sercia, P. (2019). Relationship between daily training loads and perceptions of wellness in Canadian University volleyball athletes. *Journal of Australian Strength and Conditioning*, 27(7), 8-12.
- Roy, X., Caya, O., Charron, J., Comtois, A.S., & Sercia, P. (2020). Using global and differential ratings of perceived exertion to measure internal training load in university volleyball players. *Journal of Australian Strength and Conditioning*, 28(2), 6-13.
- Ryan, R.M., & Deci, E.L. (2000). Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *American Psychologist*, 55(1), 68-78. <https://doi.org/10.1037/0003-066X.55.1.68>

- Ryan, S., Kempton, T., & Coutts, A.J. (2021). Data reduction approaches to athlete monitoring in professional Australian football. *International Journal of Sports Physiology and Performance*, 16(1), 59-65. <https://doi.org/10.1123/IJSPP.2020-0083>
- Sanders, G.J., Boos, B., Shipley, F., Scheadler, C.M., & Peacock, C.A. (2018). An accelerometer-based training load analysis to assess volleyball performance. *Journal of Exercise and Nutrition*, 1(1), 2-6. <https://doi.org/10.53520/jen2018.1034>
- Sattler, T. (2021). Prospektivna analiza poškodb in spremljanje obremenitev v odbojki s tehnologijo My Vert. [Prospective analysis of sports injury and load monitoring in volleyball team.] *Sport: Revija Za Teoreticna in Prakticna Vprasanja Sporta*, 69(3/4), 175-182.
- Saw, A.E., Main, L.C., & Gastin, P.B. (2016). Monitoring the athlete training response: Subjective self-reported measures trump commonly used objective measures: A systematic review. *British Journal of Sports Medicine*, 50(5), 281-291. <https://doi.org/10.1136/bjsports-2015-094758>
- Scott, B.R., Duthie, G.M., Thornton, H.R., & Dascombe, B.J. (2016). Training monitoring for resistance exercise: Theory and applications. *Sports Medicine*, 46(5), 687-698. <https://doi.org/10.1007/s40279-015-0454-0>
- Sheppard, J.M., & Newton, R.U. (2012). Long-term training adaptations in elite male volleyball players. *Journal of Strength and Conditioning Research*, 26(8), 2180-2184. <https://doi.org/10.1519/JSC.0b013e31823c429a>
- Sheppard, J.M., Nolan, E., & Newton, R.U. (2012). Changes in strength and power qualities over two years in volleyball players transitioning from junior to senior national team. *Journal of Strength and Conditioning Research*, 26(1), 152-157. <https://doi.org/10.1519/JSC.0b013e31821e4d5b>
- Skazalski, C., Whiteley, R., & Bahr, R. (2018). High jump demands in professional volleyball-large variability exists between players and player positions. *Scandinavian Journal of Medicine and Science in Sports*, 28(11), 2293-2298. <https://doi.org/10.1111/sms.13255>
- Sole, C.J., Kavanaugh, A.A., & Stone, M.H. (2017). Injuries in collegiate women's volleyball: A four-year retrospective analysis. *Sports*, 5(2), 26. <https://doi.org/10.3390/sports5020026>
- Souglis, A., Bogdanis, G.C., Giannopoulou, I., Papadopoulos, C., & Apostolidis, N. (2015). Comparison of inflammatory responses and muscle damage indices following a soccer, basketball, volleyball and handball game at an elite competitive level. *Research in Sports Medicine*, 23(1), 59-72. <https://doi.org/10.1080/15438627.2014.975814>
- Sousa, D. (2014). Validation in qualitative research: General aspects and specificities of the descriptive phenomenological method. *Qualitative Research in Psychology*, 11(2), 211-227. <https://doi.org/10.1080/14780887.2013.853855>
- Tavares, F., Simões, M., Matos, B., Smith, T.B., & Driller, M. (2018). Wellness, muscle soreness and neuromuscular performance during a training week in volleyball athletes. *The Journal of Sports Medicine and Physical Fitness*, 58(12), 1852-1858. <https://doi.org/10.23736/S0022-4707.17.07818-5>
- Taylor, J.B., Barnes, H.C., Gombatto, S.P., Greenwood, D., & Ford, K.R. (2022). Quantifying external load and injury occurrence in women's collegiate volleyball players across a competitive season. *Journal of Strength and Conditioning Research*, 36(3), 805-812. <https://doi.org/10.1519/JSC.0000000000004212>
- Taylor, J.B., Kantor, J.L., Hockenjos, T.J., Barnes, H.C., & Dischiavi, S.L. (2019). Jump load and landing patterns of collegiate female volleyball players during practice and competition. *Journal of Sports Medicine and Physical Fitness*, 59(11), 1892-1896. <https://doi.org/10.23736/S0022-4707.19.09650-6>
- Timoteo, T.F., Debieen, P.B., Miloski, B.D., Werneck, F.Z., Gabbett, T., & Bara Filho, M.G. (2021). Influence of workload and recovery on injuries in elite male volleyball players. *Journal of Strength and Conditioning Research*, 35(3), 791-796. <https://doi.org/10.1519/JSC.0000000000002754>
- Timoteo, T.F., Seixas, M.B., Almeida Falci, M.F., Debieen, P.B., Miloski, B.D., Miranda, R., & Bara Filho, M.G. (2017). Impact of consecutive games on workload, state of recovery and well-being of professional volleyball players. *Journal of Exercise Physiology Online*, 20(3), 130-140.
- Ungureanu, A.N., Brustio, P.R., Boccia, G., Rainoldi, A., & Lupo, C. (2021). Effects of pre-session well-being perception on internal training load in female volleyball players. *International Journal of Sports Physiology and Performance*, 16(5), 622-627. <https://doi.org/10.1123/ijssp.2020-0387>
- Ungureanu, A.N., Lupo, C., Boccia, G., & Brustio, P.R. (2021). Internal training load affects day-after-pretraining perceived fatigue in female volleyball players. *International Journal of Sports Physiology and Performance*, 16(12), 1844-1850. <https://doi.org/10.1123/ijssp.2020-0829>
- van der Does, H.T.D., Sanne Brink, M., Ardi Otter, R.T., Visscher, C., & Plechelmus Marie Lemmink, K.A. (2017). Injury risk is increased by changes in perceived recovery of team sport players. *Clinical Journal of Sport Medicine*, 27(1), 46-51. <https://doi.org/10.1097/JSM.0000000000000306>
- Vavassori, R., Moreno, M.P., & Ureña Espa, A. (2023). The perception of volleyball student-athletes: Evaluation of well-being, sport workload, players' response, and academic demands. *Healthcare (Switzerland)*, 11(11), 1538. <https://doi.org/10.3390/healthcare11111538>
- Visnes, H., & Bahr, R. (2013). Training volume and body composition as risk factors for developing jumper's knee among young elite volleyball players. *Scandinavian Journal of Medicine and Science in Sports*, 23(5), 607-613. <https://doi.org/10.1111/j.1600-0838.2011.01430.x>
- Vlantes, T.G., & Readdy, T. (2017). Using microsensor technology to quantify match demands in collegiate women's volleyball. *Journal of Strength and Conditioning Research*, 31(12), 3266-3278. <https://doi.org/10.1519/JSC.0000000000002208>

- Wolfe, H., Poole, K., Villasante Tezanos, A.G., English, R., Uhl, T.L., Tezanos, A.G.V., English, R., & Uhl, T.L. (2019). Volleyball overhead swing volume and injury frequency over the course of a season. *International Journal of Sports Physical Therapy*, 14(1), 88-96. <https://doi.org/10.26603/ijsp20190088>
- Xu, J., Wang, S., Liu, Z., Wang, Y., Wang, Y., & Dang, Y. (2023). Information theory on change to reflection. In *SpringerBriefs in Computer Science*. Springer. https://doi.org/10.1007/978-981-19-9929-1_1
- Xue, Y. (2017). Study on the application of Trimp theory based on heart rate in university volleyball training monitoring. *Agro Food Industry Hi-Tech*, 28(1), 3520-3523.
- Zahavi, D. (2020). The practice of phenomenology: The case of Max van Manen. *Nursing Philosophy*, 21(2), 1-9. <https://doi.org/10.1111/nup.12276>
- Zhou, X. (2021). Research on monitoring volleyball players' competition load based on intelligent tracking technology. *Microprocessors and Microsystems*, 82, 103881. <https://doi.org/10.1016/j.micpro.2021.103881>

Submitted: September 29, 2023

Accepted: June 5, 2024

Published Online First: June 27, 2024

Correspondence to:

Roberto Vavassori, Ph.D. candidate

University of Granada, 18011 Granada, Spain

Tel.: +34619422185

E-mail: roberto_vavassori@hotmail.com

Acknowledgements

This review will be part of a PhD thesis of RV, titled Effect of exam periods and competition on wellness and readiness in high school and university volleyball players. University of Granada, Spain.

Funding details: The authors did not receive any specific grant for this research from any funding agency in the public, commercial, or not-for-profit sectors.

Declaration of interest statement: RV, MPMA and AU declare that they have no conflict of interest.

Data availability: The datasets generated and/or analyzed during the current review are available from the corresponding author on a reasonable request.